arXiv:1304.7427v1 [math.AG] 28 Apr 2013

AN ALGORITHM TO COMPUTE AUTOMORPHISM GROUPS OF $K3$ SURFACES

ICHIRO SHIMADA

Abstract. Let X be an algebraic complex $K3$ surface or a supersingular $K3$ surface in odd characteristic. We present an algorithm by which, under certain assumptions on X, we can calculate a finite set of generators of the image of the natural homomorphism from the automorphism group of X to the orthogonal group of its Néron-Severi lattice.

1. Introduction

The automorphism group $\mathrm{Aut}(X)$ of an algebraic $K3$ surface X is an interesting object. Suppose that X is defined over the complex number field $\mathbb{C}$, or is supersingular in odd characteristic. Then, thanks to the Torelli-type theorem due to Piatetski-Shapiro and Shafarevich [24] and Ogus [22], [23], we can study $\mathrm{Aut}(X)$ by the Néron-Severi lattice S_X of X. We denote by $\mathrm{O}(S_X)$ the orthogonal group of S_X. Then we have a natural homomorphism

$$\mathrm{Aut}(X) \to \mathrm{O}(S_X).$$

It is known that this homomorphism has only a finite kernel. Using the reduction theory for arithmetic subgroups of $\mathrm{O}(S_X)$, Sterk [30] and Lieblich and Maulik [18] proved that $\mathrm{Aut}(X)$ is finitely generated. The Néron-Severi lattices for which $\mathrm{Aut}(X)$ is finite were classified by Nikulin [20], [21] and Vinberg [32]. On the other hand, when $\mathrm{Aut}(X)$ is infinite, it is in general a difficult problem to give a set of generators.

The purpose of this paper is to present a lattice-theoretic algorithm (Algorithm 5.14) that calculates a finite set of generators of the image of the homomorphism $\mathrm{Aut}(X) \to \mathrm{O}(S_X)$ under certain assumptions on X. These assumptions are described in Section 6.3. Theoretically, this algorithm can be applied to a wide class of $K3$ surfaces. In practice, however, the computation seems to be intractable in general. Nevertheless, in Section 7, we demonstrate this algorithm on some complex elliptic $K3$ surfaces with Picard number 3.

Example 1.1. Let X be a complex $K3$ surface with Picard number 3 such that a Gram matrix of S_X with respect to a certain basis of S_X is equal to

$$M = \begin{bmatrix} 0 & 1 & 0 \\ 1 & -2 & 0 \\ 0 & 0 & -24 \end{bmatrix},$$

2000 *Mathematics Subject Classification.* 14J28, 14J50, 14Q10.

Partially supported by JSPS Grant-in-Aid for Challenging Exploratory Research No.23654012 and JSPS Grants-in-Aid for Scientific Research (C) No.25400042 .

that the vector $a \in S_X$ written as $[2, 1, 0]$ with respect to this basis is ample, and that the period is sufficiently general in $T_X \otimes \mathbb{C}$, where T_X is the transcendental lattice of X. We let $\mathrm{O}(S_X)$ act on S_X from the right so that

$$\mathrm{O}(S_X) = \{\, g \in \mathrm{GL}_3(\mathbb{Z}) \mid g\, M\, {}^t g = M \,\}.$$

Then the natural homomorphism $\mathrm{Aut}(X) \to \mathrm{O}(S_X)$ is injective, and its image is generated by the following matrices:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \quad \begin{bmatrix} 1 & 0 & 0 \\ 12 & 1 & -1 \\ 24 & 0 & -1 \end{bmatrix}, \quad \begin{bmatrix} 37 & 12 & -5 \\ 36 & 13 & -5 \\ 360 & 120 & -49 \end{bmatrix}, \quad \begin{bmatrix} 97 & 48 & -14 \\ 0 & 1 & 0 \\ 672 & 336 & -97 \end{bmatrix}.$$

Moreover we can show that $\mathrm{Aut}(X)$ acts on the set of (-2)-curves on X transitively, and that there exists only one Jacobian fibration on X modulo $\mathrm{Aut}(X)$.

Even when the computation is too heavy to complete in a practical time, this algorithm yields many interesting automorphisms of a $K3$ surface.

Our method is based on a generalization of Borcherds' method [4], [5]. Suppose that an even hyperbolic lattice S of rank < 26 is primitively embedded into the even unimodular hyperbolic lattice $II := \mathrm{II}_{1,25}$ of rank 26. Let $\mathcal{P}_{II} \subset II \otimes \mathbb{R}$ be a positive cone of II. Then the hyperplanes $(r)^\perp$ perpendicular to the vectors $r \in II$ with $r^2 = -2$ decompose $\mathcal{P}_{II}$ into a union of closed chambers, which we call $\mathcal{R}^*_{II}$*-chambers*. Let $\mathcal{P}_S$ be the positive cone of $S \otimes \mathbb{R}$ that is contained in $\mathcal{P}_{II}$ by the fixed embedding. Then the tessellation of $\mathcal{P}_{II}$ by the $\mathcal{R}^*_{II}$-chambers induces a tessellation of $\mathcal{P}_S$. We call the chambers of this induced tessellation $\mathcal{R}^*_{II|S}$*-chambers*. Borcherds [4, Lemma 5.1] proved that, if the orthogonal complement R of S in II is a root lattice, then the $\mathcal{R}^*_{II|S}$-chambers are congruent to each other by the action of $\mathrm{O}^+(S)$, where $\mathrm{O}^+(S) \subset \mathrm{O}(S)$ is the subgroup of index 2 consisting of elements that leave $\mathcal{P}_S$ invariant. Using this fact, one can obtain a set of generators of $\mathrm{O}^+(S)$ for some lattices S. Kondo [16] applied this method to the Néron-Severi lattice of a generic Jacobian Kummer surface Y, and described $\mathrm{Aut}(Y)$. Since then, automorphism groups of several $K3$ surfaces have been determined by this method (Dolgachev-Kondo [12], Kondo-Keum [15], Dolgachev-Keum [13], Kondo-Shimada [17]). The classical two examples of Vinberg [31] can be also treated by this method.

We extend Borcherds' method to the situation in which II is not necessarily of rank 26 and R is not necessarily a root lattice. In this case, there are several $\mathrm{O}^+(S)$-congruence classes of $\mathcal{R}^*_{II|S}$-chambers. Starting from an $\mathcal{R}^*_{II|S}$-chamber, we compute $\mathcal{R}^*_{II|S}$-chambers adjacent to the $\mathcal{R}^*_{II|S}$-chambers obtained so far until no new $\mathrm{O}^+(S)$-congruence classes appear. By the reduction theory for arithmetic subgroups of $\mathrm{O}(S)$, we can show that this algorithm terminates and gives a finite set of generators of $\mathrm{O}^+(S)$. By means of the Torelli-type theorem, this algorithm is modified for the computation of a set of generators of the image of $\mathrm{Aut}(X) \to \mathrm{O}(S_X)$.

This paper is organized as follows. In Section 2, we fix notions and notation about lattices and hyperbolic spaces. In Section 3, we introduce the notion of chamber decomposition of a positive cone of a hyperbolic lattice, and present some algorithms about chambers. In Section 4, we review Conway theory [7] on the structure of $\mathcal{R}^*_{II}$-chambers in the even hyperbolic unimodular lattice $II = \mathrm{II}_{1,n-1}$

of rank $n = 10, 18$ or 26. Then, in Section 5, we introduce the notion of $\mathcal{R}^*_{II|S}$-chambers associated with a primitive embedding $S \hookrightarrow II$ of an even hyperbolic lattice S, and present our main algorithm (Algorithm 5.14). In Section 6, we explain how to apply Algorithm 5.14 to the study of the automorphism groups of $K3$ surfaces. In Section 7, we demonstrate Algorithm 5.14 on some $K3$ surfaces X with Picard number 3 by embedding S_X into $\mathrm{II}_{1,9}$ primitively. In Section 8, we give a brief account of examples whose computation seems to be intractable.

In this paper, *Aut* denotes the automorphism group of a lattice theoretic object, while Aut denotes the automorphism group of a $K3$ surface.

For the actual computation, we used the `C` library `gmp` [14].

2. Preliminaries on lattices and hyperbolic spaces

Let L be a free $\mathbb{Z}$-module of finite rank. We say that a submodule M of L is *primitive* if L/M is torsion-free, and that $v \in L$ is *primitive* if so is the submodule $\langle v \rangle := \mathbb{Z}v \subset L$.

Let L be a lattice; that is, a free $\mathbb{Z}$-module of finite rank with a non-degenerate symmetric bilinear form $\langle\ ,\ \rangle_L : L \times L \to \mathbb{Z}$. The subscript L in $\langle\ ,\ \rangle_L$ is omitted if no confusion will occur. The symmetric bilinear form on $L \otimes \mathbb{R}$ obtained from $\langle\ ,\ \rangle$ is also denoted by $\langle\ ,\ \rangle$. We denote by $L^\vee := \mathrm{Hom}(L, \mathbb{Z})$ the *dual lattice of* L, which is naturally embedded in $L \otimes \mathbb{Q}$. We say that L is *unimodular* if $L = L^\vee$ holds. The *norm* $\langle v, v \rangle$ of $v \in L \otimes \mathbb{R}$ is denoted by v^2. We let the *orthogonal group* $\mathrm{O}(L)$ of L act on L from the right, and write the action of $g \in \mathrm{O}(L)$ on $v \in L \otimes \mathbb{R}$ by $v \mapsto v^g$.

A lattice L is said to be *even* if $x^2 \in 2\mathbb{Z}$ holds for any $x \in L$. For an even lattice L, we put

$$\mathcal{R}_L := \{\ v \in L \ \mid\ v^2 = -2\ \}.$$

Elements of $\mathcal{R}_L$ are called (-2)*-vectors*. Each $r \in \mathcal{R}_L$ defines the *reflection*

$$s_r : x \mapsto x + \langle x, r \rangle r,$$

which is an element of $\mathrm{O}(L)$. We denote by $W(L)$ the subgroup of $\mathrm{O}(L)$ generated by all the reflections s_r with respect to (-2)-vectors $r \in \mathcal{R}_L$.

A lattice L of rank $n > 0$ is said to be *negative-definite* if the signature of the real quadratic space $L \otimes \mathbb{R}$ is $(0, n)$. A negative-definite lattice L is said to be a *root lattice* if L is generated by $\mathcal{R}_L$.

A lattice L of rank $n > 1$ is said to be *hyperbolic* if the signature of $L \otimes \mathbb{R}$ is $(1, n-1)$. Let L be a hyperbolic lattice. Then a *positive cone of* L is one of the two connected components of $\{x \in L \otimes \mathbb{R} \mid x^2 > 0\}$. The closure of a positive cone $\mathcal{P}_L$ in $L \otimes \mathbb{R}$ is denoted by $\overline{\mathcal{P}}_L$. We denote by $\overline{\mathcal{P}}_L^{\mathbb{Q}}$ the convex hull of $\overline{\mathcal{P}}_L \cap (L \otimes \mathbb{Q})$. The stabilizer subgroup in $\mathrm{O}(L)$ of a positive cone is denoted by $\mathrm{O}^+(L)$. We have $\mathrm{O}(L) = \mathrm{O}^+(L) \times \{\pm 1\}$. Note that $W(L)$ is contained in $\mathrm{O}^+(L)$.

The following algorithms will be used frequently in this paper.

Algorithm 2.1. Let Q be a positive-definite symmetric matrix of size n with rational entries, ℓ a column vector of length n with rational entries, and c a rational number. Then we can calculate the list of all row vectors $x \in \mathbb{Z}^n$ satisfying

$$x\, Q\, {}^t x + 2\, x\, \ell + c = 0$$

by the method described in [28, Section 4] or [29, Section 3.1]. ∎

Algorithm 2.2. Let L be a hyperbolic lattice, let v be a vector of $L \otimes \mathbb{Q}$ with $v^2 > 0$, let α be a rational number, and let d be an integer. Then the finite set

$$\{ \, x \in L \mid \langle x, v \rangle = \alpha, \;\; \langle x, x \rangle = d \, \}$$

can be calculated by the method described in [29, Sections 3.2]. ∎

Algorithm 2.3. Let L be a hyperbolic lattice, let v, h be vectors of $L \otimes \mathbb{Q}$ such that

$$\langle v, h \rangle > 0, \;\; \langle h, h \rangle > 0, \;\; \langle v, v \rangle > 0,$$

and let d be a negative integer. Then the finite set

$$\{ \, x \in L \mid \langle v, x \rangle < 0, \;\; \langle h, x \rangle > 0, \;\; \langle x, x \rangle = d \, \}$$

can be calculated by the method described in [29, Sections 3.3]. ∎

Let L be a hyperbolic lattice of rank $m+1$, and let $\mathcal{P}_L$ be a positive cone of L. The multiplicative group $\mathbb{R}_{>0}$ of positive real numbers acts on $\overline{\mathcal{P}}_L \setminus \{0\}$ by scalar multiplication. We put

$$\overline{\mathbb{H}}_L := (\overline{\mathcal{P}}_L \setminus \{0\})/\mathbb{R}_{>0}, \quad \overline{\mathbb{H}}_L^{\mathbb{Q}} := (\overline{\mathcal{P}}_L^{\mathbb{Q}} \setminus \{0\})/\mathbb{R}_{>0}, \quad \mathbb{H}_L := \mathcal{P}_L/\mathbb{R}_{>0},$$
$$\partial\,\overline{\mathbb{H}}_L := \overline{\mathbb{H}}_L \setminus \mathbb{H}_L, \quad \partial\,\overline{\mathbb{H}}_L^{\mathbb{Q}} := \overline{\mathbb{H}}_L^{\mathbb{Q}} \setminus \mathbb{H}_L,$$

and denote by $\pi_L : \overline{\mathcal{P}}_L \setminus \{0\} \to \overline{\mathbb{H}}_L$ the natural projection. The space $\mathbb{H}_L$ is naturally endowed with a structure of the hyperbolic m-space. A point of $\partial\,\overline{\mathbb{H}}_L^{\mathbb{Q}}$ is called a *rational boundary point*. For simplicity, when we are given a subset T of $\overline{\mathcal{P}}_L$, we denote by $\pi_L(T) \subset \overline{\mathbb{H}}_L$ the image of $T \cap (\overline{\mathcal{P}}_L \setminus \{0\})$ by π_L. For example, we have $\partial\,\overline{\mathbb{H}}_L^{\mathbb{Q}} = \pi_L(\overline{\mathcal{P}}_L \cap L)$. A subset K of $\mathbb{H}_L$ is said to be a *linear subspace of* $\mathbb{H}_L$ if there exists a linear subspace $\tilde{K}$ of $L \otimes \mathbb{R}$ such that $K = \pi_L(\mathcal{P}_L \cap \tilde{K})$ holds. Let b be a point of $\partial\,\overline{\mathbb{H}}_L$. A linear subsapce K of $\mathbb{H}_L$ is said to *pass through b at infinity* if the linear subspace $\tilde{K}$ of $L \otimes \mathbb{R}$ such that $K = \pi_L(\mathcal{P}_L \cap \tilde{K})$ contains a non-zero vector v such that $b = \pi_L(v)$.

We denote by $\mathcal{H}_L$ the set of hyperplanes of $\mathcal{P}_L$. We put

$$\mathcal{N}_L := \{ \, v \in L \otimes \mathbb{R} \mid v^2 < 0 \, \},$$

and, for $v \in L \otimes \mathbb{R}$, we put

$$[v]^{\perp} := \{ \, x \in L \otimes \mathbb{R} \mid \langle x, v \rangle = 0 \, \} \quad \text{and} \quad (v)^{\perp} := [v]^{\perp} \cap \mathcal{P}_L.$$

Since $[v]^{\perp}$ intersects $\mathcal{P}_L$ if and only if $v^2 < 0$, the map $v \mapsto (v)^{\perp}$ induces a bijection from $\mathcal{N}_L/\mathbb{R}^{\times}$ to $\mathcal{H}_L$. For $v \in \mathcal{N}_L$, we obtain a hyperplane $\pi_L((v)^{\perp})$ of the hyperbolic space $\mathbb{H}_L$.

We recall some properties of *horospheres*. For details, see Ratcliffe [25, Chapter 4]. Let b be a point of $\partial\,\overline{\mathbb{H}}_L$. Consider the upper halfspace model

$$\{ \, (z_1, \dots, z_m) \in \mathbb{R}^m \mid z_1 > 0 \, \} \tag{2.1}$$

of $\mathbb{H}_L$ such that b corresponds to the infinite point given by $z_1 = \infty$. Then every horosphere HS_b with the base b is defined by $z_1 = \gamma$ with some positive real constant γ. Therefore, as a Riemannian submanifold of $\mathbb{H}_L$, every horosphere is isomorphic to a Euclidean affine space. If K is a linear subspace of $\mathbb{H}_L$ that passes through b at infinity, then $K \cap HS_b$ is an affine subspace of any horosphere HS_b with the base b. We say that a horosphere defined by $z_1 = \gamma$ is *smaller* than a horosphere defined by $z_1 = \gamma'$ if $\gamma > \gamma'$.

Let f be a non-zero vector in $\overline{\mathcal{P}}_L$ with $f^2 = 0$, and we put $b := \pi_L(f) \in \partial\,\overline{\mathbb{H}}_L$. Let v be an element of $\mathcal{N}_L$. We can measure the distance between the hyperplane $\pi_L((v)^\perp)$ of $\mathbb{H}_L$ and b by the non-negative real number $-\langle v, f\rangle^2/v^2$. For a horosphere HS_b with the base b, we put

$$c(f, HS_b) := \sup \left\{ \, -\frac{\langle v, f\rangle^2}{v^2} \;\middle|\; v \in \mathcal{N}_L, \;\; \pi_L((v)^\perp) \cap HS_b \neq \emptyset \right\}.$$

Lemma 2.4. *Let f and b be as above. Then, for any positive real number c', there exists a horosphere HS_b with the base b such that, for any horosphere HS'_b with the base b smaller than HS_b, we have $c(f, HS'_b) < c'$.*

Proof. We choose linear coordinates $(x_0, x_1, \dots, x_m)$ of $L \otimes \mathbb{R}$ such that the quadratic form $x \mapsto x^2$ is given by

$$(x_0, x_1, \dots, x_m) \mapsto x_0^2 - x_1^2 - x_2^2 - \cdots - x_m^2,$$

that $\mathcal{P}_L$ is contained in the halfspace $x_0 > 0$, and that $f = (1, 1, 0, \dots, 0)$. We put

$$\mathbf{H}_L := \{ \, (x_0, x_1, \dots, x_m) \in L \otimes \mathbb{R} \;\mid\; x_0 > 0, \;\; x_0^2 - x_1^2 - x_2^2 - \cdots - x_m^2 = 1 \, \},$$

which is isomorphic to $\mathbb{H}_L$ as a metric space by $\pi_L|\mathbf{H}_L : \mathbf{H}_L \xrightarrow{\sim} \mathbb{H}_L$. We also put

$$\mathbf{B}_L := \{ \, (y_1, \dots, y_m) \in \mathbb{R}^m \;\mid\; y_1^2 + \cdots + y_m^2 < 1 \, \}.$$

The *conformal ball model of* $\mathbb{H}_L$ is then obtained by the isomorphism $\beta : \mathbf{H}_L \xrightarrow{\sim} \mathbf{B}_L$ given by

$$\beta(x_0, x_1, \dots, x_m) := \left(\frac{x_1}{1 + x_0}, \dots, \frac{x_m}{1 + x_0} \right),$$

by which the point $b \in \partial\,\overline{\mathbb{H}}_L$ corresponds to $(1, 0, \dots, 0) \in \partial\,\overline{\mathbf{B}}_L$, where $\overline{\mathbf{B}}_L$ is the closure of $\mathbf{B}_L$ in $\mathbb{R}^m$. The horospheres with the base b are given in $\mathbf{B}_L$ by

$$\mathbf{HS}(\alpha) : (y_1 - \alpha)^2 + y_2^2 + \cdots + y_m^2 - (1 - \alpha)^2 = 0$$

by some real numbers α with $0 < \alpha < 1$, and the parameter α tends to 1 when the horosphere $\mathbf{HS}(\alpha)$ becomes smaller. Let $v = (v_0, v_1, \dots, v_m)$ be a vector in $\mathcal{N}_L$. Suppose that there exists a point $x = (x_0, x_1, \dots, x_m) \in \mathbf{H}_L$ such that $x \in (v)^\perp$ and that $\beta(x) \in \mathbf{HS}(\alpha)$. It is enough to show that

$$-\frac{\langle v, f\rangle^2}{v^2} \leq \frac{(1 - \alpha)^2}{\alpha^2}. \tag{2.2}$$

If $\langle v, f\rangle = 0$, then (2.2) holds. Suppose that $\langle v, f\rangle \neq 0$. Replacing v by $-v$ if necessary, we can assume that $\langle v, f\rangle = v_0 - v_1$ is positive. Since $\beta(x) \in \mathbf{HS}(\alpha)$, we have

$$(x_1 - (1 + x_0)\alpha)^2 + x_2^2 + \cdots + x_m^2 - (1 - \alpha)^2 (1 + x_0)^2 = 0.$$

Since $x_2^2 + \cdots + x_m^2 = x_0^2 - x_1^2 - 1$ and $1 + x_0 \neq 0$, we obtain

$$\alpha(x_0 - x_1) - (1 - \alpha) = 0.$$

Using $\langle v, x\rangle = v_0 x_0 - v_1 x_1 - \cdots - v_m x_m = 0$, we obtain

$$x_0 = \frac{\alpha(v_2 x_2 + \cdots + v_m x_m) - (1 - \alpha) v_1}{\alpha(v_0 - v_1)}, \quad x_1 = \frac{\alpha(v_2 x_2 + \cdots + v_m x_m) - (1 - \alpha) v_0}{\alpha(v_0 - v_1)}.$$

Combining these with $x_0^2 - x_1^2 - x_2^2 - \cdots - x_m^2 - 1 = 0$ and using $v_0 \neq v_1$, we get

$$g_2(x_2) + \cdots + g_m(x_m) + \alpha^2(v_0 - v_1) + (1-\alpha)^2(v_0 + v_1) = 0, \tag{2.3}$$

where

$$g_i(t) = \alpha^2(v_0 - v_1)t^2 - 2(\alpha - \alpha^2)v_i t = \alpha^2(v_0 - v_1)\left(t - \frac{(1-\alpha)v_i}{\alpha(v_0 - v_1)}\right)^2 - \frac{(1-\alpha)^2 v_i^2}{v_0 - v_1}.$$

Since $v_0 - v_1 > 0$ and $g_i(t) \ge -(1-\alpha)^2 v_i^2/(v_0 - v_1)$, we obtain from (2.3) that

$$(1-\alpha)^2 v^2 + \alpha^2 \langle f, v\rangle^2 \le 0,$$

which is equivalent to (2.2). □

3. Chamber decomposition

Let L be an even hyperbolic lattice. For a subset Δ of $\mathcal{N}_L$, we put

$$\Sigma_L(\Delta) := \{\, x \in L \otimes \mathbb{R} \;\mid\; \langle x, v\rangle \ge 0 \;\text{ for all }\; v \in \Delta \,\}.$$

Definition 3.1. A closed subset D of $\mathcal{P}_L$ is called a *chamber* if its interior D° is non-empty and there exists a subset Δ of $\mathcal{N}_L$ such that $D = \Sigma_L(\Delta) \cap \mathcal{P}_L$ holds. Let D be a chamber. A hyperplane $(v)^\perp \in \mathcal{H}_L$ is called a *wall of* D if $(v)^\perp \cap D^\circ$ is empty and $(v)^\perp \cap D$ contains a non-empty open subset of $(v)^\perp$. We denote by $\overline{D}$ the closure of D in $\overline{\mathcal{P}}_L$.

Definition 3.2. Let $\mathcal{F} \subset \mathcal{H}_L$ be a locally finite family of hyperplanes in $\mathcal{P}_L$. Then the closure in $\mathcal{P}_L$ of each connected component of

$$\mathcal{P}_L \setminus \bigcup_{(v)^\perp \in \mathcal{F}} (v)^\perp$$

is a chamber, which we call an $\mathcal{F}$*-chamber.*

By definition, every wall of an $\mathcal{F}$-chamber is an element of $\mathcal{F}$. If D and D' are distinct $\mathcal{F}$-chambers, then $D^\circ \cap D' = \emptyset$ holds.

Definition 3.3. Let D be an $\mathcal{F}$-chamber, and let $(v)^\perp \in \mathcal{F}$ is a wall of D. Then there exists a unique $\mathcal{F}$-chamber D' such that $D' \neq D$ and that $D \cap D' \cap (v)^\perp$ contains a non-empty open subset of $(v)^\perp$. We say that D' is *adjacent to* D *along the wall* $(v)^\perp$.

We fix a subgroup G of $\mathrm{O}^+(L)$ with finite index. We assume the following:

- [G] There exists an algorithm by which we can determine, for a given $g \in \mathrm{O}^+(L)$, whether $g \in G$ or not.

Let $\mathcal{V}$ be a subset of $\mathcal{N}_L \cap L^\vee$, and consider the family of hyperplanes

$$\mathcal{V}^* := \{\, (v)^\perp \;\mid\; v \in \mathcal{V} \,\}$$

in $\mathcal{P}_L$. We assume that $\mathcal{V}$ has the following properties:

- [V1] There exists $c \in \mathbb{R}$ such that, for any $v \in \mathcal{V}$, we have $-v^2 < c$.
- [V2] The set $\mathcal{V}$ is invariant under the action of G on $\mathcal{N}_L \cap L^\vee$.

Then we can consider $\mathcal{V}^*$-chambers by the following:

Lemma 3.4. *The family of hyperplanes $\mathcal{V}^*$ is locally finite in $\mathcal{P}_L$.*

Proof. Since $\{v^2 \mid v \in L^\vee\}$ is discrete in $\mathbb{R}$, the set $\{v^2 \mid v \in \mathcal{V}\}$ is finite by the property [V1] and $\mathcal{V} \subset \mathcal{N}_L \cap L^\vee$. Since $L^\vee$ is discrete in $L \otimes \mathbb{R}$, we obtain the proof of Lemma 3.4. □

Definition 3.5. Let D be a $\mathcal{V}^*$-chamber. A subset Δ of $\mathcal{N}_L \cap L^\vee$ is called a *defining set of* D if $D = \Sigma_L(\Delta) \cap \mathcal{P}_L$ holds. (Note that Δ need not be contained in $\mathcal{V}$.) A defining set Δ of D is said to be *minimal* if the following hold:

- For any $v \in \Delta$, the hyperplane $(v)^\perp$ is a wall of D, and
- if v and v' are distinct vectors of Δ, then $(v)^\perp \neq (v')^\perp$.

We define two types of minimal defining sets, each of which is unique for a given $\mathcal{V}^*$-chamber D. The one is called the *$\mathcal{V}$-minimal defining set*, denoted by $\Delta_{\mathcal{V}}(D)$, and characterized by the following property:

- $\Delta_{\mathcal{V}}(D) \subset \mathcal{V}$, and if $v \in \Delta_{\mathcal{V}}(D)$, then $\alpha v \notin \mathcal{V}$ for any $\alpha \in \mathbb{R}$ with $0 < \alpha < 1$.

The other is called the *primitively minimal defining set*, denoted by $\Delta_{L^\vee}(D)$, and characterized by the following property:

- Every $v \in \Delta_{L^\vee}(D)$ is primitive in $L^\vee$.

Let D be a $\mathcal{V}^*$-chamber. Then D^g is also a $\mathcal{V}^*$-chamber for any $g \in G$ by the property [V2]. For a $\mathcal{V}^*$-chamber D, we put

$$Aut_G(D) := \{\, g \in G \mid D^g = D \,\}.$$

Let D and D' be $\mathcal{V}^*$-chambers. We say that D and D' are *G-congruent* if there exists $g \in G$ such that $D' = D^g$. The following simple observation is the key point of our method:

$$\text{If } g \in G \text{ satisfies } D^g \cap D'^\circ \neq \emptyset, \text{ then } D^g = D'. \tag{3.1}$$

In particular, if $g \in G$ satisfies $D^g \cap D^\circ \neq \emptyset$, then $g \in Aut_G(D)$.

Example 3.6. The subset $\mathcal{R}_L$ of $\mathcal{N}_L \cap L^\vee$ has the properties [V1] and [V2] for $G = \mathrm{O}^+(L)$. Let D be an $\mathcal{R}_L^*$-chamber. Then D is a fundamental domain of the action of $W(L)$ on $\mathcal{P}_L$, and $W(L)$ is generated by the reflections s_r, where r runs through the $\mathcal{R}_L$-minimal defining set $\Delta_{\mathcal{R}_L}(D)$ of D. Moreover, any two $\mathcal{R}_L^*$-chambers are $\mathrm{O}^+(L)$-congruent, and $\mathrm{O}^+(L)$ is isomorphic to the semi-direct product $W(L) \rtimes Aut_{\mathrm{O}^+(L)}(D)$.

We consider the following properties of $\mathcal{V}$:

[V3] Any $\mathcal{V}^*$-chamber has a *finite* defining set.

[V4] For any $\mathcal{V}^*$-chamber D, the set $\pi_L(\overline{D}) \cap \partial\,\overline{\mathbb{H}}_L$ is contained in $\partial\,\overline{\mathbb{H}}_L^{\mathbb{Q}}$.

The main results of this section are the following:

Theorem 3.7. *Suppose that $\mathcal{V}$ satisfies* [V1]-[V4]. *Then there exist only a finite number of G-congruence classes of $\mathcal{V}^*$-chambers.*

Theorem 3.8. *Suppose that $\mathcal{V}$ satisfies* [V1]-[V4]. *Then the group $Aut_G(D)$ is finite for any $\mathcal{V}^*$-chamber D.*

3.1. **Proof of Theorem 3.7.** We assume that $\mathcal{V}$ satisfies [V1]-[V4].

Definition 3.9. A subset Π of $\overline{\mathcal{P}}_L^{\mathbb{Q}}$ is called a *rational polyhedral cone* if there exist a finite number of non-zero vectors $v_1, \dots, v_n \in \overline{\mathcal{P}} \cap L$ such that

$$\Pi = \mathbb{R}_{\geq 0} v_1 + \cdots + \mathbb{R}_{\geq 0} v_n.$$

Recall that G is assumed to be of finite index in $\mathrm{O}^+(L)$. We have the following result from the reduction theory of arithmetic subgroups of $\mathrm{O}(L)$ (see Ash et al. [2, Chapter II, Section 4] and Sterk [30]).

Theorem 3.10. *There exist a finite number of rational polyhedral cones* $\Pi_1, \dots, \Pi_N$ *in* $\overline{\mathcal{P}}_L^{\mathbb{Q}}$ *such that* $\mathcal{P}_L$ *is equal to*

$$\bigcup_{g\in G} \bigcup_{i=1}^{N} (\Pi_i^g \cap \mathcal{P}_L).$$

Therefore Theorem 3.7 follows from the following:

Proposition 3.11. *Let* Π *be a rational polyhedral cone in* $\overline{\mathcal{P}}_L^{\mathbb{Q}}$. *Then the number of* $\mathcal{V}^*$*-chambers that intersect* $\Pi \cap \mathcal{P}_L$ *is finite.*

For the proof of Proposition 3.11, we need two corollaries of Lemma 2.4.

Let b be a point of $\partial\,\overline{\mathbb{H}}_L$. A *closed horoball* HB_b with the base b is a subset of $\mathbb{H}_L$ defined by $z_1 \ge \gamma$ with some positive real constant γ in the upper halfspace model (2.1) with b at $z_1 = \infty$. Let ∂HB_b be the horosphere defined by $z_1 = \gamma$. The map $\rho_b : HB_b \to \partial HB_b$ defined by

$$\rho_b(z_1, z_2, \dots, z_m) := (\gamma, z_2, \dots, z_m)$$

is called the *natural projection.* Let $b = \pi_L(f) \in \partial\,\overline{\mathbb{H}}_L^{\mathbb{Q}}$ be a rational boundary point, where f is a non-zero vector in $\overline{\mathcal{P}}_L \cap L$ with $f^2 = 0$. We put

$$\mathcal{V}_b := \{\ v \in \mathcal{V} \ \mid\ \langle v, f\rangle = 0\ \}.$$

If $v \in \mathcal{V}$ satisfies $v \notin \mathcal{V}_b$, then we have $\langle v, f\rangle^2 \ge 1$ because $f \in L$ and $\mathcal{V} \subset L^\vee$. By the property [V1], there exists a positive real number δ_b such that

$$\delta_b < -\langle v, f\rangle^2/v^2 \quad \text{for any } v \in \mathcal{V} \setminus \mathcal{V}_b.$$

Therefore we obtain the following corollary of Lemma 2.4:

Corollary 3.12. *Let* b *be a rational boundary point, and let* HB_b *be a sufficiently small closed horoball with the base* b.

(1) *Let* v *be an element of* $\mathcal{V}$. *Then the hyperplane* $\pi_L((v)^\perp)$ *of* $\mathbb{H}_L$ *intersects* HB_b *if and only if* $\pi_L((v)^\perp)$ *passes through* b *at infinity.*

(2) *Let* D *be a* $\mathcal{V}^*$*-chamber. If* $b \notin \pi_L(\overline{D})$, *then* $\pi_L(D) \cap HB_b$ *is empty, while if* $b \in \pi_L(\overline{D})$, *then* $\pi_L(D) \cap HB_b = \rho_b^{-1}(\pi_L(D) \cap \partial HB_b)$ *holds.*

We regard ∂HB_b as a Euclidean affine space. Then the family of affine hyperplanes

$$\{\ \pi_L((v)^\perp) \cap \partial HB_b \ \mid\ v \in \mathcal{V}_b\ \}$$

of ∂HB_b is locally finite, because $\mathcal{V}^*$ is locally finite in $\mathcal{P}_L$ and $\pi_L^{-1}(\partial HB_b) \subset \mathcal{P}_L$. Therefore we obtain the following:

Corollary 3.13. *Let* b *and* HB_b *be as in Corollary 3.12, and let* J *be a compact subset of* ∂HB_b. *Then the number of* $\mathcal{V}^*$*-chambers that intersect the subset* $\pi_L^{-1}(\rho_b^{-1}(J))$ *of* $\mathcal{P}_L$ *is finite.*

Proof of Proposition 3.11. Let $v_1, \dots, v_n \in \overline{\mathcal{P}}_L \cap L$ be non-zero vectors such that Π is equal to $\mathbb{R}_{\ge 0}v_1 + \cdots + \mathbb{R}_{\ge 0}v_n$. We number $v_1, \dots, v_n$ in such a way that

$$v_1^2 = \cdots = v_k^2 = 0 \quad \text{and} \quad v_{k+1}^2 > 0,\ \dots,\ v_n^2 > 0.$$

Let b_i be the rational boundary point $\pi_L(v_i)$ for $i = 1, \dots, k$, and let HB_i be a sufficiently small closed horoball with the base b_i. The natural projection is denoted by $\rho_i : HB_i \to \partial HB_i$. We put $HB_i^\circ := HB_i \setminus \partial HB_i$. Since $\pi_L(\Pi) \cap \partial\, \overline{\mathbb{H}}_L = \{b_1, \dots, b_k\}$, we see that

$$\pi_L(\Pi)' := \pi_L(\Pi) \setminus \bigcup_{i=1}^{k} (HB_i^\circ \cap \pi_L(\Pi))$$

is a compact subset of $\mathbb{H}_L$. Therefore the number of $\mathcal{V}^*$-chambers D such that $\pi_L(D)$ intersects $\pi_L(\Pi)'$ is finite. For $j \neq i$, let $p_{i,j} \in \partial HB_i$ denote the intersection point of ∂HB_i and the geodesic line in $\mathbb{H}_L$ passing through b_i at infinity and passing through $\pi_L(v_j)$ possibly at infinity. Then the convex hull J_i of these points $p_{i,j}$ with $j \neq i$ in the Euclidean affine space ∂HB_i is compact and satisfies $HB_i \cap \pi_L(\Pi) = \rho_i^{-1}(J_i)$. Consequently, the number of $\mathcal{V}^*$-chambers D such that $\pi_L(D)$ intersects $HB_i \cap \pi_L(\Pi)$ is finite by Corollary 3.13. □

3.2. **Proof of Theorem 3.8.** We continue to assume that $\mathcal{V}$ satisfies [V1]-[V4].

Lemma 3.14. *Let Δ be a subset of $\mathcal{N}_L \cap L^\vee$ such that $D = \Sigma_L(\Delta) \cap \mathcal{P}_L$ is a $\mathcal{V}^*$-chamber. Then $\Sigma_L(\Delta)$ is contained in $\overline{\mathcal{P}}_L$.*

Proof. Note that $\Sigma_L(\Delta)$ is a closed convex subset of $L \otimes \mathbb{R}$. Suppose that there exists $x_0 \in \Sigma_L(\Delta)$ such that $x_0 \notin \overline{\mathcal{P}}_L$. Let U be a non-empty open subset of D°. Then, for any $y \in U$, the line segment $\overline{x_0 y}$ of $L \otimes \mathbb{R}$ connecting x_0 and y is contained in $\Sigma_L(\Delta)$, and $\overline{x_0 y} \cap \mathcal{P}_L$ is contained in D. Hence the intersection point $z(y)$ of $\overline{x_0 y}$ and $\{x \in \overline{\mathcal{P}}_L \mid x^2 = 0\}$ belongs to the closure $\overline{D}$ of D in $\overline{\mathcal{P}}_L$. Since U is open, the subset $\pi_L(\{z(y) \mid y \in U\})$ of $\pi_L(\overline{D}) \cap \partial\, \overline{\mathbb{H}}_L$ has uncountably many points, which contradicts the property [V4] of $\mathcal{V}$. □

Lemma 3.15. *Let D be a $\mathcal{V}^*$-chamber. Then any defining set Δ of D spans $L \otimes \mathbb{R}$.*

Proof. Let V be the linear subspace of $L \otimes \mathbb{R}$ spanned by Δ. The orthogonal complement $V^\perp$ of V in $L \otimes \mathbb{R}$ is contained in $\Sigma_L(\Delta)$ by definition and hence $V^\perp \subset \overline{\mathcal{P}}_L$ by Lemma 3.14. This holds only when $V^\perp = 0$. □

Lemma 3.16. *Suppose that a defining set Δ_1 of a $\mathcal{V}^*$-chamber D satisfies the following: If $v, v' \in \Delta_1$ are distinct, then $(v)^\perp \neq (v')^\perp$ holds. Let v be an element of Δ_1. (1) The hyperplane $(v)^\perp$ is a wall of D if and only if $\Sigma_L(\Delta_1) \neq \Sigma_L(\Delta_1 \setminus \{v\})$. (2) If $\Delta_1 \setminus \{v\}$ does not span $L \otimes \mathbb{R}$, then $(v)^\perp$ is a wall of D.*

Proof. The 'only if' part of (1) is obvious by the assumption on Δ_1. Conversely, suppose that $\Sigma_L(\Delta_1) \neq \Sigma_L(\Delta_1 \setminus \{v\})$. Then there exists $x_0 \in L \otimes \mathbb{R}$ such that $\langle v, x_0 \rangle < 0$ and $x_0 \in \Sigma_L(\Delta_1 \setminus \{v\})$. Assume that $(v)^\perp$ is not a wall of D. Then D is equal to $\Sigma_L(\Delta_1 \setminus \{v\}) \cap \mathcal{P}_L$, and hence $\Sigma_L(\Delta_1 \setminus \{v\})$ is contained in $\overline{\mathcal{P}}_L$ by Lemma 3.14. In particular, we have $x_0 \in \overline{\mathcal{P}}_L$. Let y_0 be a point of D°. Then the intersection point z_0 of the line segment $\overline{x_0 y_0}$ and $(v)^\perp$ satisfies $z_0^2 > 0$, $\langle v, z_0 \rangle = 0$ and $\langle v', z_0 \rangle > 0$ for any v' in $\Delta_1 \setminus \{v\}$. Since $\mathcal{V}^*$ is locally finite in $\mathcal{P}_L$, these mean that a sufficiently small open neighborhood of z_0 in $(v)^\perp$ is contained in D. In anyway, $(v)^\perp$ is a wall of D. Thus (1) is proved. If $(v)^\perp$ is not a wall of D, then $\Delta_1 \setminus \{v\}$ is also a defining set of D. Hence (2) follows from Lemma 3.15. □

Algorithm 3.17. Let Δ be a *finite* defining set of a $\mathcal{V}^*$-chamber D. This algorithm calculates the primitively minimal defining set $\Delta_{L^\vee}(D)$ of D.

Step 0. We set $\Delta_1 := \{\}$ and $\Delta_2 := \{\}$.

Step 1. For each element $v \in \Delta$, we calculate the maximal positive integer a_v such that $v/a_v \in L^\vee$, and append v/a_v to Δ_1. Then we have $D = \Sigma_L(\Delta_1) \cap \mathcal{P}_L$, and moreover, if $v, v' \in \Delta_1$ are distinct, then $(v)^\perp \neq (v')^\perp$ holds.

Step 2. For each $v \in \Delta_1$, we carry out the following computation. Suppose that $\Delta_1 \setminus \{v\}$ does not span $L \otimes \mathbb{R}$. Then $(v)^\perp$ is a wall of D by Lemma 3.16, and we append v to Δ_2. Suppose that $\Delta_1 \setminus \{v\}$ spans $L \otimes \mathbb{R}$. Then we can solve the following problem of linear programming on the vector space $L \otimes \mathbb{Q}$:

$$\begin{cases} \text{minimize} & \langle v, x\rangle \\ \text{subject to} & \langle v', x\rangle \geq 0 \;\; \text{for all} \;\; v' \in \Delta_1 \setminus \{v\}. \end{cases} \tag{3.2}$$

(See, for example, Chvátal [6] for the algorithms of linear programming.) Note that the solution is either 0 or unbounded to $-\infty$. If the solution is 0, then $(v)^\perp$ is not a wall of D by Lemma 3.16. Suppose that the solution is unbounded to $-\infty$. Then there exists $x_0 \in L \otimes \mathbb{R}$ such that $\langle v, x_0\rangle < 0$ and $x_0 \in \Sigma_L(\Delta_1 \setminus \{v\})$. Hence $(v)^\perp$ is a wall of D by Lemma 3.16, and we append v to Δ_2.

Step 3. We then output Δ_2 as $\Delta_{L^\vee}(D)$. ∎

Remark that, for any $\mathcal{V}^*$-chamber D, the minimal defining set $\Delta_{L^\vee}(D)$ is finite by the property [V3] of $\mathcal{V}$ and Algorithm 3.17.

Algorithm 3.18. Suppose that the primitively minimal defining set $\Delta_{L^\vee}(D)$ of a $\mathcal{V}^*$-chamber D is given. This algorithm calculates all elements of $Aut_G(D)$. Let $\Delta_{L^\vee}(D)^l$ denote the set of ordered l-tuples of distinct elements of $\Delta_{L^\vee}(D)$, where $l := \operatorname{rank} L$. By Lemma 3.15, there exists $[v_1, \dots, v_l] \in \Delta_{L^\vee}(D)^l$ that forms a basis of $L \otimes \mathbb{Q}$. We set $A := \{\}$. For each $[v'_1, \dots, v'_l] \in \Delta_{L^\vee}(D)^l$, we calculate the linear transformation g of $L \otimes \mathbb{Q}$ such that

$$v_i^g = v'_i \qquad (i = 1, \dots, l).$$

If g belongs to G and induces a permutation of $\Delta_{L^\vee}(D)$, then we append g to A. When this calculation is done for all $[v'_1, \dots, v'_l] \in \Delta_{L^\vee}(D)^l$, the set A is equal to $Aut_G(D)$. ∎

Proof of Theorem 3.8. Since $\Delta_{L^\vee}(D)$ is finite, Algorithm 3.18 terminates in finite steps. □

Algorithm 3.19. Let D and D' be $\mathcal{V}^*$-chambers. Suppose that $\Delta_{L^\vee}(D)$ and $\Delta_{L^\vee}(D')$ are given. This algorithm determines whether D is G-congruent to D' or not. We fix an element $[v_1, \dots, v_l]$ of $\Delta_{L^\vee}(D)^l$ that forms a basis of $L \otimes \mathbb{Q}$. For each $[v''_1, \dots, v''_l] \in \Delta_{L^\vee}(D')^l$, we calculate the linear transformation g of $L \otimes \mathbb{Q}$ that satisfies $v_i^g = v''_i$ for $i = 1, \dots, l$. If g belongs to G and induces a bijection from $\Delta_{L^\vee}(D)$ to $\Delta_{L^\vee}(D')$, then D and D' are G-congruent. If no such $[v''_1, \dots, v''_l]$ are found, then D and D' are not G-congruent. ∎

Remark 3.20. Suppose that we somehow know a point $p \in D^\circ$. By (3.1), we see that an element $g \in G$ is contained in $Aut_G(D)$ if and only if $p^g \in D$, and $g \in G$ induces an isomorphism from D to D' if and only if $p^g \in D'$.

4. Conway theory

Let n be 10, 18 or 26. Throughout this section, we denote by II an even *unimodular* hyperbolic lattice $\mathrm{II}_{1,n-1}$ of rank n. Note that II exists and is unique up to isomorphisms (see, for example, Serre [27, Chapter V]). We fix a positive cone $\mathcal{P}_{II}$ of II. Conway [7] ([11, Chapter 27]) described the structure of $\mathcal{R}^*_{II}$-chambers.

Definition 4.1. Let $\mathcal{D}$ be an $\mathcal{R}^*_{II}$-chamber. We say that a vector $w \in II$ is a *Weyl vector of* $\mathcal{D}$ if the $\mathcal{R}_{II}$-minimal defining set $\Delta_{\mathcal{R}_{II}}(\mathcal{D})$ of $\mathcal{D}$ is given by

$$\Delta_{\mathcal{R}_{II}}(\mathcal{D}) = \{ \, r \in \mathcal{R}_{II} \;\mid\; \langle w, r \rangle = 1 \, \}.$$

If a Weyl vector of an $\mathcal{R}^*_{II}$-chamber $\mathcal{D}$ exists, then it is unique, because, as will be shown below, $\Delta_{\mathcal{R}_{II}}(\mathcal{D})$ spans $II \otimes \mathbb{R}$. If w is a Weyl vector of $\mathcal{D}$, then w^g is a Weyl vector of $\mathcal{D}^g$ for any $g \in \mathrm{O}^+(II)$. Since any two $\mathcal{R}^*_{II}$-chambers are $\mathrm{O}^+(II)$-congruent by Example 3.6, a Weyl vector of a single $\mathcal{R}^*_{II}$-chamber gives Weyl vectors of all $\mathcal{R}^*_{II}$-chambers via the action of $\mathrm{O}^+(II)$.

Theorem 4.2 (Conway [7])**.** *For any $\mathcal{R}^*_{II}$-chamber $\mathcal{D}$, there exists a Weyl vector $w \in II$ of $\mathcal{D}$. We have*

$$w^2 = \begin{cases} 1240 & \text{if } n = 10 \\ 620 & \text{if } n = 18 \\ 0 & \text{if } n = 26 \end{cases} \qquad \text{and} \qquad |\Delta_{\mathcal{R}_{II}}(\mathcal{D})| = \begin{cases} 10 & \text{if } n = 10 \\ 19 & \text{if } n = 18 \\ \infty & \text{if } n = 26. \end{cases}$$

We give an explicit description of the Weyl vectors, and prove the following:

Proposition 4.3. *If $\mathcal{D}$ is an $\mathcal{R}^*_{II}$-chamber, then $\pi_{II}(\overline{\mathcal{D}}) \cap \partial\, \overline{\mathbb{H}}_{II}$ is contained in $\partial\, \overline{\mathbb{H}}^{\mathbb{Q}}_{II}$.*

For a non-zero vector v of $II \otimes \mathbb{R}$, we denote by

$$[v]_+ := \{ \, av \;\mid\; a \in \mathbb{R}_{\geq 0} \, \}$$

the half-line generated by v. Let Δ be a *finite* defining set of an $\mathcal{R}^*_{II}$-chamber. We say that $[v]_+$ is an *extremal edge* of $\Sigma_{II}(\Delta)$ if the following hold:

- $[v]_+$ is contained in $\Sigma_{II}(\Delta)$, and
- there exist linearly independent vectors $r_1, \dots, r_{n-1} \in \Delta$ such that $\langle v, r_i \rangle = 0$ holds for $i = 1, \dots, n-1$.

We can make the complete list of extremal edges of $\Sigma_{II}(\Delta)$ by executing the following calculation to all non-ordered $(n-1)$-tuples $\{r_1, \dots, r_{n-1}\}$ of Δ. We determine whether $r_1, \dots, r_{n-1}$ are linearly independent or not. If they are linearly independent, we calculate a non-zero vector v that satisfies $\langle v, r_i \rangle = 0$ for $i = 1, \dots, n-1$. Then we calculate $\langle v, r \rangle$ for all $r \in \Delta$. If they are all non-negative, then $[v]_+$ is an extremal edge of $\Sigma_{II}(\Delta)$, while if they are all non-positive, then $[-v]_+$ is an extremal edge. Otherwise, the linearly independent set $\{r_1, \dots, r_{n-1}\}$ does not yield an extremal edge.

In the following, we denote by U the even hyperbolic lattice of rank 2 with a fixed basis f_U, z_U, with respect to which the Gram matrix is

$$\begin{bmatrix} 0 & 1 \\ 1 & -2 \end{bmatrix},$$

and by $f_U^\vee, z_U^\vee$ the basis of U dual to f_U, z_U. Let E_8 denote the (negative-definite) root lattice of type E_8 with the standard basis $e_1, \dots, e_8$, whose Coxeter graph is



We denote by $e_1^\vee, \dots, e_8^\vee$ the basis of E_8 dual to $e_1, \dots, e_8$. We put

$$\theta := 3e_1 + 2e_2 + 4e_3 + 6e_4 + 5e_5 + 4e_6 + 3e_7 + 2e_8.$$

Finally, we denote by Λ the *negative-definite* Leech lattice; that is, Λ is an even unimodular negative-definite lattice of rank 24 such that $\mathcal{R}_\Lambda = \emptyset$.

4.1. **The case where $n = 10$.** We put $I\!I := U \oplus E_8$, and choose $\mathcal{P}_{I\!I}$ in such a way that $2f_U + z_U \in \mathcal{P}_{I\!I}$. Then the vector

$$w_0 := 30f_U^\vee + z_U^\vee + e_1^\vee + \cdots + e_8^\vee$$

is a Weyl vector of an $\mathcal{R}_{I\!I}^*$-chamber $\mathcal{D}_0$. We have

$$\Delta_{\mathcal{R}_{I\!I}}(\mathcal{D}_0) = \{z_U, e_1, \dots, e_8, f_U - \theta\}.$$

We confirm that there exist 10 extremal edges $[v_0]_+, \dots, [v_9]_+$ of $\Sigma_{I\!I}(\Delta_{\mathcal{R}_{I\!I}}(\mathcal{D}_0))$. Each v_i satisfies $v_i^2 \geq 0$, and hence we have $\Sigma_{I\!I}(\Delta_{\mathcal{R}_{I\!I}}(\mathcal{D}_0)) \subset \overline{\mathcal{P}}_{I\!I}$. Moreover there exists exactly one extremal edge among them such that $v_i^2 = 0$, which is equal to $[f_U]_+$. Thus we have

$$\pi_{I\!I}(\overline{\mathcal{D}}_0) \cap \partial\, \overline{\mathbb{H}}_{I\!I} = \{\pi_{I\!I}(f_U)\} \subset \partial\, \overline{\mathbb{H}}_{I\!I}^{\mathbb{Q}}.$$

4.2. **The case where $n = 18$.** We put $I\!I := U \oplus E_8 \oplus E_8$, and choose $\mathcal{P}_{I\!I}$ in such a way that $2f_U + z_U \in \mathcal{P}_{I\!I}$. Let $e'_1, \dots, e'_8$ be the basis of the second E_8 with the same Coxeter graph as $e_1, \dots, e_8$. Then the vector

$$w_0 := 30f_U^\vee + z_U^\vee + e_1^\vee + \cdots + e_8^\vee + e_1'^\vee + \cdots + e_8'^\vee$$

is a Weyl vector of an $\mathcal{R}_{I\!I}^*$-chamber $\mathcal{D}_0$. We have

$$\Delta_{\mathcal{R}_{I\!I}}(\mathcal{D}_0) = \{z_U, e_1, \dots, e_8, e'_1, \dots, e'_8, f_U - \theta, f_U - \theta'\},$$

where θ' is defined in the same way as θ with e_i replaced by e'_i. There exist 82 extremal edges $[v_0]_+, \dots, [v_{81}]_+$ of $\Sigma_{I\!I}(\Delta_{\mathcal{R}_{I\!I}}(\mathcal{D}_0))$. Each v_i satisfies $v_i^2 \geq 0$, and hence we have $\Sigma_{I\!I}(\Delta_{\mathcal{R}_{I\!I}}(\mathcal{D}_0)) \subset \overline{\mathcal{P}}_{I\!I}$. Moreover there exist exactly two extremal edges among them such that $v_i^2 = 0$, which are given by $[f_U]_+$ and $[v_1]_+$, where

$$v_1 = e_1 + e_3 + e'_1 + e'_3 + 2(z_U + (f_U - \theta) + (f_U - \theta') + e_4 + \cdots + e_8 + e'_4 + \cdots + e'_8).$$

Hence we have

$$\pi_{I\!I}(\overline{\mathcal{D}}_0) \cap \partial\, \overline{\mathbb{H}}_{I\!I} = \{\pi_{I\!I}(f_U), \pi_{I\!I}(v_1)\} \subset \partial\, \overline{\mathbb{H}}_{I\!I}^{\mathbb{Q}}.$$

4.3. **The case where $n = 26$.** We put $I\!I := U \oplus \Lambda$, and choose $\mathcal{P}_{I\!I}$ in such a way that $2f_U + z_U \in \mathcal{P}_{I\!I}$.

Theorem 4.4 (Conway and Sloane [9])**.** *A vector $w \in I\!I$ is a Weyl vector if and only if w is a non-zero primitive vector of norm 0 contained in $\overline{\mathcal{P}}_{I\!I}$ such that $\langle w\rangle^\perp/\langle w\rangle$ is isomorphic to Λ, where $\langle w\rangle^\perp$ is the orthogonal complement of the primitive submodule $\langle w\rangle$ of $I\!I$.*

Therefore $w_0 := f_U$ is a Weyl vector of an $\mathcal{R}_{I\!I}^*$-chamber $\mathcal{D}_0$. For simplicity, we denote vectors of $I\!I \otimes \mathbb{R} = (U \oplus \Lambda) \otimes \mathbb{R}$ by

$$[s, t, y] := sf_U + tz_U + y, \quad \text{where} \quad s, t \in \mathbb{R}, \ \ y \in \Lambda \otimes \mathbb{R}.$$

Then we have

$$\Delta_{\mathcal{R}_{I\!I}}(\mathcal{D}_0) = \{\ r_\lambda \ \mid\ \lambda \in \Lambda\ \}, \quad \text{where} \quad r_\lambda := \left[-\lambda^2/2, 1, \lambda\right].$$

Conway, Parker and Sloane [8] ([11, Chapter 23]) proved that the covering radius of the Leech lattice is $\sqrt{2}$ (see also Borcherds [3]). A point c of $\Lambda \otimes \mathbb{R}$ is called a *deep hole* if $-(c-\lambda)^2 \geq 2$ holds for any $\lambda \in \Lambda$. In [4], Borcherds observed the following:

Lemma 4.5. *Let b be a point of $\pi_{II}(\overline{\mathcal{D}}_0) \cap \partial\, \overline{\mathbb{H}}_{II}$. Then either $b = \pi_{II}(w_0)$ or there exists a deep hole c such that*

$$b = \pi_{II}(v_c), \quad \textit{where} \quad v_c := \left[-c^2/2 + 1, 1, c\right].$$

In particular, the set $\pi_{II}(\overline{\mathcal{D}}_0) \cap \partial\, \overline{\mathbb{H}}_{II}$ is contained in $\partial\, \overline{\mathbb{H}}_{II}^{\mathbb{Q}}$.

Proof. Note that $\overline{\mathcal{P}}_{II}$ is contained in $\{x \in II \otimes \mathbb{R} \mid \langle w_0, x\rangle \geq 0\}$, and that $\overline{\mathcal{P}}_{II} \cap [w_0]^{\perp}$ is the half-line $[w_0]_+$. Suppose that $b = \pi_{II}(u)$, where $u = [s, t, y]$ is a non-zero vector of norm 0 in $\overline{\mathcal{D}}_0$. Since $u \in \overline{\mathcal{P}}_{II}$, we have $t = \langle w_0, u\rangle \geq 0$, and $t = 0$ holds if and only if $[w_0]_+ = [u]_+$. Hence $t = 0$ implies $b = \pi_{II}(w_0)$. Suppose that $t \neq 0$. We can assume that $t = 1$. Since $u^2 = 0$, we have $2s - 2 + y^2 = 0$. Since $u \in \overline{\mathcal{D}}_0$, we have

$$\langle u, r_\lambda\rangle = -\frac{(y-\lambda)^2}{2} - 1 \geq 0 \;\text{ for any }\; \lambda \in \Lambda.$$

Therefore $y \in \Lambda \otimes \mathbb{R}$ is a deep hole c and $u = v_c$ holds. Since every deep hole belongs to $\Lambda \otimes \mathbb{Q}$ by Conway, Parker and Sloane [8] or Conway and Sloane [10] ([11, Chapters 23 and 24]), we have $\pi_{II}(u) \in \partial\, \overline{\mathbb{H}}_{II}^{\mathbb{Q}}$. $\square$

5. Generalized Borcherds' method

Suppose that S is an even hyperbolic lattice with a fixed positive cone $\mathcal{P}_S$, and let G be a subgroup of $\mathrm{O}^+(S)$ with finite index satisfying [G] in Section 3 with L replaced by S. We present an algorithm that calculates a set of generators of $Aut_G(E)$ for a given $\mathcal{R}_S^*$-chamber E under the assumptions [SG1], [SG2] and [SG3] below.

We recall the definition of the discriminant form of an even lattice. See Nikulin [19] for details. For an even lattice L, the *discriminant group* $A_L := L^\vee/L$ is equipped with a non-degenerate quadratic form

$$q_L : A_L \to \mathbb{Q}/2\mathbb{Z}, \qquad x \bmod L \mapsto x^2 \bmod 2\mathbb{Z},$$

which is called the *discriminant form of L*. Let $\mathrm{O}(q_L)$ denote the group of automorphisms of (A_L, q_L). We have a natural homomorphism

$$\eta_L : \mathrm{O}(L) \to \mathrm{O}(q_L).$$

Let n be 10, 18 or 26. As in Section 4, we denote by II an even unimodular hyperbolic lattice of rank n. We assume that

[SG1] the rank of S is smaller that n, and S is primitively embedded into II.

Let R denote the orthogonal complement of S in II. Then R is an even negative-definite lattice. We assume that

[SG2] if $n = 26$, the lattice R cannot be embedded into the Leech lattice Λ.

For example, if $\mathcal{R}_R$ is non-empty, then [SG2] is satisfied, because $\mathcal{R}_\Lambda = \emptyset$.

We denote by

$$x \mapsto x_S \quad \text{and} \quad x \mapsto x_R$$

the orthogonal projections from $II \otimes \mathbb{R}$ to $S \otimes \mathbb{R}$ and $R \otimes \mathbb{R}$, respectively. Since II is contained in $S^\vee \oplus R^\vee$, the images of II by these projections are contained in $S^\vee$ and

$R^\vee$, respectively. Since $I\!I$ is unimodular, the result of Nikulin [19, Proposition 1.6.1] implies that the subgroup $I\!I/(S \oplus R)$ of $A_S \oplus A_R$ is the graph of an isomorphism

$$\delta : (A_S, q_S) \xrightarrow{\sim} (A_R, -q_R).$$

Let $\delta_* : \mathrm{O}(q_S) \xrightarrow{\sim} \mathrm{O}(q_R)$ denote the isomorphism induced by δ. We assume that the subgroup G of $\mathrm{O}^+(S)$ satisfies

[SG3] $\delta_*(\eta_S(G)) \subset \operatorname{Im} \eta_R$.

For example, if G is contained in $\eta_S^{-1}(\{\pm 1\})$, then [SG3] is satisfied.

Let $\mathcal{P}_{I\!I}$ be the positive cone of $I\!I$ that contains the fixed positive cone $\mathcal{P}_S$ of S. Let r be an element of $\mathcal{R}_{I\!I}$. Then the hyperplane $(r)^\perp \in \mathcal{H}_{I\!I}$ of $\mathcal{P}_{I\!I}$ intersects $\mathcal{P}_S$ if and only if $r_S^2 < 0$ holds, and in this case, the hyperplane $(r_S)^\perp \in \mathcal{H}_S$ of $\mathcal{P}_S$ is equal to the intersection $\mathcal{P}_S \cap (r)^\perp$. We put

$$\mathcal{R}_{I\!I|S} := \{\, r_S \mid r \in \mathcal{R}_{I\!I},\ r_S^2 < 0 \,\},$$

and show that the subset $\mathcal{R}_{I\!I|S}$ of $\mathcal{N}_S \cap S^\vee$ has the properties [V1]-[V4] given in Section 3 with L replaced by S.

Proposition 5.1. *If $v \in \mathcal{R}_{I\!I|S}$, then $-v^2 \le 2$.*

Proof. Since R is negative-definite and $r_S^2 + r_R^2 = -2$ holds for any $r \in \mathcal{R}_{I\!I}$, we have $-2 \le v^2$ for any $v = r_S \in \mathcal{R}_{I\!I|S}$. □

Therefore the family of hyperplanes

$$\begin{aligned} \mathcal{R}^*_{I\!I|S} &= \{\, (r_S)^\perp \in \mathcal{H}_S \mid r \in \mathcal{R}_{I\!I},\ r_S^2 < 0 \,\} \\ &= \{\, \mathcal{P}_S \cap (r)^\perp \mid r \in \mathcal{R}_{I\!I},\ \mathcal{P}_S \cap (r)^\perp \ne \emptyset \,\} \end{aligned}$$

is locally finite in $\mathcal{P}_S$.

Proposition 5.2. *Let g be an element of G. Then there exists $\tilde{g} \in \mathrm{O}^+(I\!I)$ that leaves S invariant and is equal to g on S.*

Proof. By [SG3], there exists $h \in \mathrm{O}(R)$ such that $\delta_*(\eta_S(g)) = \eta_R(h)$. Then the action of $(\eta_S(g), \eta_R(h))$ on $A_S \oplus A_R$ preserves the graph $I\!I/(S \oplus R)$ of δ, and hence the action of (g, h) on $S^\vee \oplus R^\vee$ preserves $I\!I \subset S^\vee \oplus R^\vee$. The restriction $\tilde{g} \in \mathrm{O}(I\!I)$ of (g, h) to $I\!I$ belongs to $\mathrm{O}^+(I\!I)$, because $\tilde{g}$ leaves $\mathcal{P}_S \subset \mathcal{P}_{I\!I}$ invariant. Thus we obtain a desired lift $\tilde{g} \in \mathrm{O}^+(I\!I)$. □

Proposition 5.3. *The action of G on $\mathcal{N}_S \cap S^\vee$ leaves $\mathcal{R}_{I\!I|S}$ invariant.*

Proof. For any $v \in \mathcal{R}_{I\!I|S}$ and $g \in G$, we have $r \in \mathcal{R}_{I\!I}$ such that $v = r_S$ and a lift $\tilde{g} \in \mathrm{O}^+(I\!I)$ of g. Then $v^g = (r^{\tilde{g}})_S \in \mathcal{R}_{I\!I|S}$ holds. □

By the definition of $\mathcal{R}_{I\!I|S}$, every $\mathcal{R}^*_{I\!I|S}$-chamber D is written as

$$D = \mathcal{D} \cap \mathcal{P}_S$$

by some $\mathcal{R}^*_{I\!I}$-chamber $\mathcal{D}$.

Remark 5.4. Let $\mathcal{D}$ be an $\mathcal{R}^*_{I\!I}$-chamber. Then $\mathcal{D} \cap \mathcal{P}_S$ is an $\mathcal{R}^*_{I\!I|S}$-chamber if and only if the interior of $\mathcal{D} \cap \mathcal{P}_S$ in $S \otimes \mathbb{R}$ is non-empty.

Definition 5.5. Let D be an $\mathcal{R}^*_{I\!I|S}$-chamber, and let $\mathcal{D}$ be an $\mathcal{R}^*_{I\!I}$-chamber such that $D = \mathcal{D} \cap \mathcal{P}_S$. By abuse of terminology, we say that the Weyl vector $w \in I\!I$ of $\mathcal{D}$ is a *Weyl vector of D*.

Proposition 5.6. *Let w and w' be Weyl vectors of an $\mathcal{R}^*_{II|S}$-chamber D. Then $w_S = w'_S$ holds.*

Proof. There exists a canonical one-to-one correspondence between the set of $\mathcal{R}^*_{II}$-chambers $\mathcal{D}$ satisfying $D = \mathcal{D} \cap \mathcal{P}_S$ and the set of connected components of

$$(R \otimes \mathbb{R}) \setminus \bigcup_{\rho \in \mathcal{R}_R} [\rho]^\perp, \quad \text{where} \quad [\rho]^\perp := \{x \in R \otimes \mathbb{R} \,|\, \langle x, \rho \rangle_R = 0\}.$$

If $\mathcal{R}^*_{II}$-chambers $\mathcal{D}$ and $\mathcal{D}'$ satisfy $D = \mathcal{D} \cap \mathcal{P}_S = \mathcal{D}' \cap \mathcal{P}_S$, then there exists a sequence of reflections s_i $(i = 1, \dots, N)$ of II with respect to $\rho_i \in \mathcal{R}_R \subset \mathcal{R}_{II}$ such that their product $s_1 \dots s_N$ maps $\mathcal{D}$ to $\mathcal{D}'$. Then the Weyl vector w of $\mathcal{D}$ is mapped to the Weyl vector w' of $\mathcal{D}'$ by $s_1 \cdots s_N$. Since each (-2)-vector ρ_i is contained in R, we have $w_S = w'_S$. □

Let D be an $\mathcal{R}^*_{II|S}$-chamber, let $w \in II$ be a Weyl vector of D, and let $\mathcal{D}$ be the $\mathcal{R}^*_{II}$-chamber corresponding to w, so that we have $D = \mathcal{D} \cap \mathcal{P}_S$. Recall that $\Delta_{R_{II}}(\mathcal{D}) = \{r \in \mathcal{R}_{II} \,|\, \langle w, r \rangle_{II} = 1\}$. We put

$$\Delta_w := \{\ r \in \Delta_{R_{II}}(\mathcal{D}) \ \mid\ r_S^2 < 0\ \} \quad \text{and} \quad \mathrm{pr}_S(\Delta_w) := \{r_S \,|\, r \in \Delta_w\}. \tag{5.1}$$

Then we have $D = \Sigma_S(\mathrm{pr}_S(\Delta_w)) \cap \mathcal{P}_S$.

Remark 5.7. It is *not* true in general that, if $w \in II$ is a Weyl vector of D, then $w_S \in S^\vee$ belongs to $\overline{D}$. It is *not* true in general either that the defining set $\mathrm{pr}_S(\Delta_w)$ of D is minimal.

Proposition 5.8. *Any $\mathcal{R}^*_{II|S}$-chamber D has a finite defining set.*

Proof. Let $\mathcal{D}$ be an $\mathcal{R}^*_{II}$-chamber such that $D = \mathcal{D} \cap \mathcal{P}_S$, and let w be the Weyl vector of $\mathcal{D}$. It is enough to show that the set Δ_w defined by (5.1) is finite. First we show $w_S^2 > 0$. Note that $w^2 = w_S^2 + w_R^2$ and $w_R^2 \le 0$. Hence $w_S^2 > 0$ holds when $n = 10$ or $n = 18$ by Theorem 4.2. Suppose that $n = 26$ and $w_S^2 = 0$. Then $w_R = 0$ and $w = w_S$ hold. Therefore $\langle w \rangle^\perp / \langle w \rangle$ would contain R, which contradicts Theorem 4.4 and the assumption [SG2].

We denote by d_R the order of A_R. Then $d_R v^2 \in \mathbb{Z}$ and $d_R^2 v^2 \in 2\mathbb{Z}$ hold for any $v \in R^\vee$. We put

$$n_R := \{\ c \in \mathbb{Q} \ \mid\ d_R c \in \mathbb{Z},\ \ d_R^2 c \in 2\mathbb{Z},\ \ -2 < c \le 0\ \}, \tag{5.2}$$

which is obviously finite. Since R is negative-definite, the set

$$R^\vee[c] := \{\ v \in R^\vee \ \mid\ v^2 = c\ \} \tag{5.3}$$

is finite for each $c \in n_R$. In particular, the set

$$a_R[c] := \{\ \langle w_R, v \rangle_R \ \mid\ v \in R^\vee[c]\ \} \tag{5.4}$$

is finite for each $c \in n_R$. Suppose that $r \in \Delta_{\mathcal{R}_{II}}(\mathcal{D})$ satisfies $r_S^2 < 0$. Note that

$$r^2 = r_S^2 + r_R^2 = -2 \quad \text{and} \quad \langle w, r \rangle_{II} = \langle w_S, r_S \rangle_S + \langle w_R, r_R \rangle_R = 1.$$

Since $r_R^2 \le 0$, we have $-2 \le r_S^2 < 0$ and $-2 < r_R^2 \le 0$. Hence $n' := r_R^2$ belongs to n_R, and r_R is an element of $R^\vee[n']$. Since $w_S^2 > 0$, the quadratic part of the inhomogeneous quadratic form $x \mapsto x^2$ on the affine hyperplane

$$\{\ x \in S \otimes \mathbb{R} \ \mid\ \langle w_S, x \rangle_S = b\ \}$$

of $S \otimes \mathbb{R}$ is negative-definite for any $b \in \mathbb{R}$. If we put $a' := \langle w_R, r_R \rangle_R \in a_R[n']$, then r_S belongs to

$$S^{\vee}[n', a'] := \{\, v \in S^{\vee} \;\mid\; \langle w_S, v \rangle_S = 1 - a', \;\; v^2 = -2 - n' \,\}. \tag{5.5}$$

Since n_R, $R^{\vee}[n']$, $a_R[n']$ and $S^{\vee}[n', a']$ are finite, Δ_w is finite. □

Proposition 5.9. *For any $\mathcal{R}^*_{I\!I|S}$-chamber D, we have $\pi_S(\overline{D}) \cap \partial\, \overline{\mathbb{H}}_S \subset \partial\, \overline{\mathbb{H}}_S^{\mathbb{Q}}$.*

Proof. Let $\mathcal{D}$ be an $\mathcal{R}^*_{I\!I}$-chamber such that $D = \mathcal{D} \cap \mathcal{P}_S$. Then we have $\overline{D} = \overline{\mathcal{D}} \cap \overline{\mathcal{P}}_S$. Since $\partial\, \overline{\mathbb{H}}_S^{\mathbb{Q}} = \partial\, \overline{\mathbb{H}}_S \cap \partial\, \overline{\mathbb{H}}_{I\!I}^{\mathbb{Q}}$ under the canonical inclusion $\overline{\mathbb{H}}_S \hookrightarrow \overline{\mathbb{H}}_{I\!I}$, the assertion follows from Proposition 4.3. □

Thus the subset $\mathcal{R}_{I\!I|S}$ of $\mathcal{N}_S \cap S^{\vee}$ has the properties [V1]-[V4]. Hence we can apply the algorithms described in Section 3 to $\mathcal{V} = \mathcal{R}_{I\!I|S}$.

Algorithm 5.10. Suppose that a Weyl vector $w \in I\!I$ of an $\mathcal{R}^*_{I\!I|S}$-chamber D is given as input. This algorithm calculates the primitively minimal defining set $\Delta_{S^{\vee}}(D)$ of D.

Step 0. We calculate $w_S \in S^{\vee}$, $w_R \in R^{\vee}$, and the set n_R defined by (5.2). We set $\Delta' := \{\}$.

Step 1. For each $c \in n_R$, we calculate $R^{\vee}[c]$ defined by (5.3) by Algorithm 2.1, and calculate $a_R[c]$ defined by (5.4).

Step 2. Using Algorithm 2.2, we calculate the finite set $S^{\vee}[n', a']$ defined by (5.5) for each pair of $n' \in n_R$ and $a' \in a_R[n']$.

Step 3. For each triple $n' \in n_R$, $v_R \in R^{\vee}[n']$ and $v_S \in S^{\vee}[n', a']$, where $a' = \langle w_R, v_R \rangle_R$, we determine whether $v_S + v_R \in S^{\vee} \oplus R^{\vee}$ belongs to $I\!I$ or not, and if $v_S + v_R \in I\!I$, then we append $r := v_S + v_R$ to Δ'.

Step 4. Note that Δ' is now equal to Δ_w defined by (5.1). We calculate the defining set $\mathrm{pr}_S(\Delta_w)$ of D.

Step 5. We then calculate $\Delta_{S^{\vee}}(D)$ from $\mathrm{pr}_S(\Delta_w)$ by Algorithm 3.17. ∎

Let w be a Weyl vector of an $\mathcal{R}^*_{I\!I|S}$-chamber D, and let v be an element of $\Delta_{S^{\vee}}(D)$. Then $(v)^{\perp}$ is a wall of D. We calculate the $\mathcal{R}^*_{I\!I|S}$-chamber that is adjacent to D along $(v)^{\perp}$. First we prepare an auxiliary algorithm.

Algorithm 5.11. Suppose that $v \in \mathcal{N}_S \cap S^{\vee}$ is given as input. We regard the hyperplane $(v)^{\perp}$ of $\mathcal{P}_S$ as a linear subspace of $\mathcal{P}_{I\!I}$ with codimension $\operatorname{rank} R + 1$. This algorithm calculates the set

$$P_v := \{\, r \in \mathcal{R}_{I\!I} \;\mid\; (v)^{\perp} \subset (r)^{\perp} \,\}. \tag{5.6}$$

We set $P := \{\}$. There exist only a finite number of $\alpha \in \mathbb{Q}$ such that $\alpha v \in S^{\vee}$ and $\alpha^2 v^2 \geq -2$. For each such rational number α and each $u \in R^{\vee}[c]$, where $c = -2 - \alpha^2 v^2$ and $R^{\vee}[c]$ defined by (5.3), we determine whether $\alpha v + u \in S^{\vee} \oplus R^{\vee}$ belongs to $I\!I$ or not, and if $\alpha v + u \in I\!I$, then we append $r := \alpha v + u \in I\!I$ to P. Then we output P as P_v. (Remark that P_v contains $\mathcal{R}_R$.) ∎

We put $[v] := \mathbb{R}v \subset L \otimes \mathbb{R}$, and consider the linear subspace

$$V := [v] \oplus (R \otimes \mathbb{R})$$

of $I\!I \otimes \mathbb{R}$. Let $x \mapsto x_V$ denote the orthogonal projection from $I\!I \otimes \mathbb{R}$ to V, and let $\langle\ ,\ \rangle_V : V \times V \to \mathbb{R}$ denote the restriction of $\langle\ ,\ \rangle_{I\!I}$ to V. Each element of P_v defined by (5.6) is contained in V. Hence $r_V = r$ holds for any $r \in P_v$. We denote by $\{\mathcal{D}_0, \dots, \mathcal{D}_m\}$ the set of $\mathcal{R}^*_{I\!I}$-chambers containing the linear subspace $(v)^\perp$ of $\mathcal{P}_{I\!I}$, and put

$$\mathcal{D}^\circ_{j,V} := \{\ x \in V \ \mid\ \langle x, r\rangle_V > 0 \ \text{ for any } \ r \in P_v \cap \Delta_{\mathcal{R}_{I\!I}}(\mathcal{D}_j)\ \}.$$

Then $\mathcal{D}_j \mapsto \mathcal{D}^\circ_{j,V}$ gives a one-to-one correspondence from $\{\mathcal{D}_0, \dots, \mathcal{D}_m\}$ to the set of connected components of

$$V \ \setminus \bigcup_{r \in P_v} [r]^\perp_V, \quad \text{where} \quad [r]^\perp_V := \{\ x \in V \ \mid\ \langle x, r\rangle_V = 0\ \}.$$

Let w_j be the Weyl vector of $\mathcal{D}_j$. By renumbering $\mathcal{D}_0, \dots, \mathcal{D}_m$, we can assume that $D = \mathcal{P}_S \cap \mathcal{D}_0$ and that w_0 is the given Weyl vector w of D. Since $\langle w_{j,V}, r\rangle_V = \langle w_j, r\rangle_{I\!I} = 1$ for any $r \in P_v \cap \Delta_{\mathcal{R}_{I\!I}}(\mathcal{D}_j)$, the vector $w_{j,V}$ belongs to $\mathcal{D}^\circ_{j,V}$. There exists $\mathcal{D}_{\mathrm{opp}}$ among $\{\mathcal{D}_0, \dots, \mathcal{D}_m\}$ such that

$$\mathcal{D}^\circ_{\mathrm{opp},V} = -\mathcal{D}^\circ_{0,V}.$$

Then $\mathcal{P}_S \cap \mathcal{D}_{\mathrm{opp}}$ is the $\mathcal{R}^*_{I\!I|S}$-chamber D' adjacent to D along $(v)^\perp$. We calculate the Weyl vector of $\mathcal{D}_{\mathrm{opp}}$. Let u be a sufficiently general vector of $I\!I \otimes \mathbb{Q}$, and let ε be a sufficiently small positive real number. Consider the oriented line segment

$$p(t) := (1-t)w_V + t(-w_V + \varepsilon u_V) \qquad (0 \le t \le 1)$$

in V from $w_V = w_{0,V} \in \mathcal{D}^\circ_{0,V}$ to $-w_V + \varepsilon u_V \in \mathcal{D}^\circ_{\mathrm{opp},V}$. Let $P'_v = \{r_1, \dots, r_N\} \subset P_v$ be a complete set of representatives of $P_v/\{\pm 1\}$. For $i = 1, \dots, N$, let t_i be the value of t such that $p(t_i) \in [r_i]^\perp_V$. Since $\langle x, r_i\rangle_{I\!I} = \langle x_V, r_i\rangle_V$ for any $x \in I\!I \otimes \mathbb{R}$, we have

$$t_i = \left(2 - \varepsilon \frac{\langle u, r_i\rangle_{I\!I}}{\langle w, r_i\rangle_{I\!I}}\right)^{-1}.$$

We put the numbering of the elements $r_1, \dots, r_N$ of P'_v so that $t_1 < \dots < t_N$ holds. Let $s_i \in \mathrm{O}^+(I\!I)$ denote the reflection with respect to r_i. Then $\mathcal{D}_0$ and $\mathcal{D}_{\mathrm{opp}}$ are related by

$$\mathcal{D}_{\mathrm{opp}} = \mathcal{D}_0^{s_1 s_2 \dots s_N}.$$

Therefore $w^{s_1 s_2 \dots s_N}$ is the Weyl vector of $\mathcal{D}_{\mathrm{opp}}$. By this consideration, we obtain the following:

Algorithm 5.12. Suppose that a Weyl vector $w \in I\!I$ of an $\mathcal{R}^*_{I\!I|S}$-chamber D and an element v of $\Delta_{S^\vee}(D)$ are given as input. This algorithm calculates a Weyl vector w' of the $\mathcal{R}^*_{I\!I|S}$-chamber D' adjacent to D along the wall $(v)^\perp$.

We calculate the set P_v by Algorithm 5.11, and choose a complete set of representatives $P'_v = \{r_1, \dots, r_N\}$ of $P_v/\{\pm 1\}$. We also choose a vector u of $I\!I \otimes \mathbb{Q}$ such that $i \ne j$ implies $\langle u, r_i\rangle_{I\!I}/\langle w, r_i\rangle_{I\!I} \ne \langle u, r_j\rangle_{I\!I}/\langle w, r_j\rangle_{I\!I}$. We sort the elements r_i of P'_v so that

$$i < j \implies \frac{\langle u, r_i\rangle_{I\!I}}{\langle w, r_i\rangle_{I\!I}} < \frac{\langle u, r_j\rangle_{I\!I}}{\langle w, r_j\rangle_{I\!I}}$$

holds. Then $w^{s_1 s_2 \dots s_N}$ is a Weyl vector of D', where $s_i \in \mathrm{O}^+(I\!I)$ is the reflection with respect to r_i. ∎

Now we present our main algorithm. Let E be an $\mathcal{R}^*_S$-chamber. Since $\mathcal{R}_S$ is contained in $\mathcal{R}_{I\!I|S}$, the $\mathcal{R}^*_S$-chamber E is a union of $\mathcal{R}^*_{I\!I|S}$-chambers. We fix an $\mathcal{R}^*_{I\!I|S}$-chamber D_0 contained in E. An *E-chain* is a finite sequence

$$D^{(0)}, D^{(1)}, \dots, D^{(N)}$$

of $\mathcal{R}^*_{II|S}$-chambers contained in E such that $D^{(a+1)}$ is adjacent to $D^{(a)}$ for each a. The *length* of an E-chain $D^{(0)}, \dots, D^{(N)}$ is defined to be N. Let D be an $\mathcal{R}^*_{II|S}$-chamber contained in E. Since E is connected, there exists an E-chain $D^{(0)}, \dots, D^{(N)}$ such that $D^{(0)} = D_0$ and $D^{(N)} = D$. The *level of* D is defined to be the minimum of the lengths of all E-chains $D^{(0)}, \dots, D^{(N)}$ such that $D^{(0)} = D_0$ and $D^{(N)} = D$.

Remark 5.13. In the actual execution of the following algorithm, we use a Weyl vector to store an $\mathcal{R}^*_{II|S}$-chamber in the computer memory. Thus, for example, the set $\mathcal{C}$ is realized as a set of vectors of II in the computer.

Algorithm 5.14. Suppose that a Weyl vector w_0 of an $\mathcal{R}^*_{II|S}$-chamber D_0 is given as input. Let E be the $\mathcal{R}^*_S$-chamber containing D_0. This algorithm calculates a finite set Γ of generators of $Aut_G(E)$, a set $\mathcal{C}$ of $\mathcal{R}^*_{II|S}$-chambers such that the union

$$F_E := \bigcup_{D_i \in \mathcal{C}} D_i$$

satisfies the property (2) in Proposition 5.17, and a set $\mathcal{B}$ of (-2)-vectors of S that satisfies the property (3) in Proposition 5.17.

Step 0. Set Γ to be $\{\}$, $\mathcal{C}_0$ to be $\{D_0\}$, and $\mathcal{B}$ to be $\{\}$.

Step 1. Calculate $\Delta_{S^\vee}(D_0)$ by Algorithm 5.10.

Step 2. Execute `adj`(0), where `adj`(ℓ) is the following procedure:

The procedure `adj`(ℓ). Suppose that, for $\lambda = 0, \dots, \ell$, a finite set $\mathcal{C}_\lambda$ of $\mathcal{R}^*_{II|S}$-chambers with the following properties has been calculated.

[C1] Each $D \in \mathcal{C}_\lambda$ is contained in E, and is of level $\le \lambda$, and

[C2] if $D, D' \in \cup_{\lambda=0}^{\ell} \mathcal{C}_\lambda$ are distinct, then D and D' are not G-congruent.

Suppose also that, for each $D \in \cup_{\lambda=0}^{\ell} \mathcal{C}_\lambda$, we have calculated $\Delta_{S^\vee}(D)$. This procedure calculates $\mathcal{C}_{\ell+1}$ and $\Delta_{S^\vee}(D)$ for each $D \in \mathcal{C}_{\ell+1}$.

We put $\mathcal{C}_\ell = \{D_{k+1}, \dots, D_{k+m}\}$, where k is the sum of the cardinalities of $\mathcal{C}_\lambda$ with $\lambda < \ell$ (we put $k = -1$ if $\ell = 0$), and m is the cardinality of $\mathcal{C}_\ell$.

1. Put $\mathcal{C}' := \{\}$ and set $l := k + m + 1$.
2. For each $D_i \in \mathcal{C}_\ell$, we make the following calculation.
   - 2-1. Calculate the finite group $Aut_G(D_i)$ from $\Delta_{S^\vee}(D_i)$ by Algorithm 3.18, and append all non-trivial elements of $Aut_G(D_i)$ to Γ.
   - 2-2. Note that $Aut_G(D_i)$ acts on $\Delta_{S^\vee}(D_i)$. Decompose $\Delta_{S^\vee}(D_i)$ into the $Aut_G(D_i)$-orbits $o_1, \dots, o_t$. Since $\mathcal{R}^*_S$ is G-invariant, the set $o^*_\nu := \{(v)^\perp \mid v \in o_\nu\}$ is either disjoint from $\mathcal{R}^*_S$ or entirely contained in $\mathcal{R}^*_S$.
   - 2-3. For each orbit o_ν such that o^*_ν is disjoint from $\mathcal{R}^*_S$, we make the following calculation.
     - 2-3-1. Choose a vector $v \in o_\nu$, and calculate a Weyl vector $w' \in II$ of the $\mathcal{R}^*_{II|S}$-chamber D' adjacent to D_i along the wall $(v)^\perp$ by Algorithm 5.12. We then calculate $\Delta_{S^\vee}(D')$ by Algorithm 5.10. Since $(v)^\perp \notin \mathcal{R}^*_S$ and $D_i \subset E$, we have $D' \subset E$. Since D_i is of level $\le \ell$, we see that D' is of level $\le \ell + 1$.
     - 2-3-2. By Algorithm 3.19, determine whether D' is G-congruent to an $\mathcal{R}^*_{II|S}$-chamber D'' contained in

$$\tilde{\mathcal{C}} := \mathcal{C}_0 \cup \dots \cup \mathcal{C}_\ell \cup \mathcal{C}' = \{D_0, D_1, \dots, D_{l-1}\}$$

or not. If ${D'}^h = D''$ for some $D'' \in \tilde{\mathcal{C}}$ and $h \in G$, then append h to $\mathit{\Gamma}$. If there exist no such $D'' \in \tilde{\mathcal{C}}$ and $h \in G$, then put $D_l := D'$, append D_l to $\mathcal{C}'$ and increment l by 1.

2-4. For each orbit o_ν such that $o_\nu^* \subset \mathcal{R}_S^*$, choose a vector $v \in o_\nu$, find a positive integer α such that $r := \alpha v \in \mathcal{R}_S$ (recall that v is primitive in $S^\vee$), and append r to $\mathcal{B}$.

3. If $\mathcal{C}' \neq \emptyset$, then put $\mathcal{C}_{\ell+1} := \mathcal{C}'$, which has the properties [C1] and [C2] above, and execute `adj`$(\ell + 1)$. If $\mathcal{C}' = \emptyset$, then put

$$\mathcal{C} := \mathcal{C}_0 \cup \mathcal{C}_1 \cup \cdots \cup \mathcal{C}_\ell,$$

and terminates.

∎

Proposition 5.15. *Algorithm 5.14 terminates.*

Proof. By construction, any two distinct $\mathcal{R}_{II|S}^*$-chambers in $\mathcal{C}_0 \cup \cdots \cup \mathcal{C}_\ell \cup \mathcal{C}'$ are not G-congruent during the calculation. Thus Proposition 5.15 follows from Theorem 3.7. □

Remark 5.16. Suppose that $n = 26$ and that R is a root lattice. By Borcherds [4, Lemma 5.1], there exists only one $\mathrm{O}^+(S)$-congruence class of $\mathcal{R}_{II|S}^*$-chambers. Hence Algorithm 5.14 terminates quickly in this case.

Proposition 5.17. (1) *The group $Aut_G(E)$ is generated by $\mathit{\Gamma}$.*

(2) *For any $v \in E$, there exists $g \in Aut_G(E)$ such that $v^g \in F_E$.*

(3) *Let r be an element of the $\mathcal{R}_S$-minimal defining set $\Delta_{\mathcal{R}_S}(E)$ of E. Then there exists $g \in Aut_G(E)$ such that $r^g \in \mathcal{B}$.*

Proof. Since each $D_i \in \mathcal{C}$ is contained in E, we have $Aut_G(D_i) \subset Aut_G(E)$ by (3.1). If $h \in G$ is appended to $\mathit{\Gamma}$ in Step 2-3-2, then we have ${D'}^h = D''$ for some $\mathcal{R}_{II|S}^*$-chambers D' and D'' contained in E, and hence $h \in Aut_G(E)$ by (3.1). Therefore the subgroup $\langle \mathit{\Gamma} \rangle$ of G generated by $\mathit{\Gamma}$ is contained in $Aut_G(E)$.

To prove Proposition 5.17, it is enough to show that, for an arbitrary $\mathcal{R}_{II|S}^*$-chamber D contained in E, there exists $\gamma \in \langle \mathit{\Gamma} \rangle$ such that $D^\gamma \in \mathcal{C}$. Indeed, let g be an arbitrary element of $Aut_G(E)$. Suppose that $(D_0^g)^\gamma = D_i \in \mathcal{C}$ for some $\gamma \in \langle \mathit{\Gamma} \rangle$. Since D_i and D_0 are G-congruent and $D_i, D_0 \in \mathcal{C}$, we have $D_0 = D_i$. Therefore $g\gamma \in Aut_G(D_0) \subset \langle \mathit{\Gamma} \rangle$ follows, and hence we have $g \in \langle \mathit{\Gamma} \rangle$. Let $v \in E$ be an arbitrary vector, and let D be an $\mathcal{R}_{II|S}^*$-chamber containing v and contained in E. If $D^\gamma = D_i \in \mathcal{C}$ for some $\gamma \in \langle \mathit{\Gamma} \rangle$, then $v^\gamma \in F_E$. Suppose that $r \in \Delta_{\mathcal{R}_S}(E)$. Then there exist an $\mathcal{R}_{II|S}^*$-chamber D contained in E and a vector $v \in \Delta_{S^\vee}(D)$ such that $r = \alpha v$ for some positive integer α. Suppose that $\gamma \in \langle \mathit{\Gamma} \rangle$ satisfies $D^\gamma = D_i \in \mathcal{C}$. Then $v^\gamma \in \Delta_{S^\vee}(D_i)$ is contained in an $Aut_G(D_i)$-orbit o_ν such that $o_\nu^* \subset \mathcal{R}_S^*$. Hence there exists $\gamma' \in Aut_G(D_i)$ such that $r^{\gamma\gamma'}$ is appended to $\mathcal{B}$ in Step 2-3-3.

We fix an $\mathcal{R}_{II|S}^*$-chamber D contained in E, and prove the existence of $\gamma \in \langle \mathit{\Gamma} \rangle$ such that $D^\gamma \in \mathcal{C}$. An E-chain $D^{(0)}, D^{(1)}, \dots, D^{(N)}$ of $\mathcal{R}_{II|S}^*$-chambers is said to be *D-admissible* if $D^{(0)}$ belongs to $\mathcal{C}$ and there exists $\gamma \in \langle \mathit{\Gamma} \rangle$ such that $D^{(N)} = D^\gamma$. Since E is connected, there exists at least one D-admissible E-chain. It is enough to show that there exists a D-admissible E-chain of length 0. We suppose that the D-admissible E-chain with *minimal* length

$$D^{(0)} = D_i,\ D^{(1)},\ \dots,\ D^{(N)} = D^\gamma \qquad (D_i \in \mathcal{C}, \gamma \in \langle \mathit{\Gamma} \rangle)$$

is of length $N>0$, and derive a contradiction. Let $v' \in \Delta_{S^\vee}(D_i)$ be the vector such that $(v')^\perp$ is the wall between $D^{(0)}=D_i$ and $D^{(1)}$. Since D_i and $D^{(1)}$ are contained in E, the $Aut_G(D_i)$-orbit $o_\nu \subset \Delta_{S^\vee}(D_i)$ containing v' satisfies $o_\nu^* \cap \mathcal{R}_S^* = \emptyset$. Let v be the vector of o_ν chosen in Step 2-3-1, and let $g \in Aut_G(D_i)$ be an element that maps v' to v. Then $D^{(1)g}$ is the $\mathcal{R}^*_{II|S}$-chamber adjacent to $D^{(0)}=D_i$ along $(v)^\perp$. Since $g \in \langle \Gamma \rangle$, the E-chain

$$D^{(0)}=D_i,\ D^{(1)g},\ \dots,\ D^{(N)g}=D^{\gamma g}$$

is D-admissible. By the minimality of N, $D^{(1)g}$ does not belong to $\mathcal{C}$. Hence, in Step 2-3-2, the $\mathcal{R}^*_{II|S}$-chamber $D^{(1)g}$ is not appended to $\mathcal{C}'$, which means that there exist $D'' \in \mathcal{C}$ and $h \in G$ such that $D^{(1)gh}=D''$. This h is appended to Γ in Step 2-3-2, and hence $\gamma gh \in \langle \Gamma \rangle$. Then

$$D^{(1)gh}=D'',\ D^{(2)gh},\ \dots,\ D^{(N)gh}=D^{\gamma gh}$$

is a D-admissible E-chain of length $N-1$, which is a contradiction. □

Remark 5.18. When Algorithm 5.14 is applicable to the case $G=\mathrm{O}^+(S)$, we see that $\mathrm{O}^+(S)$ is generated by Γ and the reflections $\{s_r \mid r \in \mathcal{B}\}$.

6. The automorphism group of a $K3$ surface

Let X be an algebraic $K3$ surface with the Néron-Severi lattice S_X of rank >1. Then S_X is an even hyperbolic lattice. Let $\mathcal{P}(X)$ denote the positive cone of S_X containing an ample class. The nef cone $\mathrm{Nef}(X)$ of X is defined by

$$\mathrm{Nef}(X) := \{\ v \in S_X \otimes \mathbb{R} \ \mid\ \langle x, [C] \rangle \ge 0 \ \text{ for any curve } C \text{ on } X\ \},$$

where $[C] \in S_X$ is the class of a curve C. It is known that $\mathrm{Nef}(X)$ is an $\mathcal{R}^*_{S_X}$-chamber, and that the $\mathcal{R}_{S_X}$-minimal defining set $\Delta_{\mathcal{R}_{S_X}}(\mathrm{Nef}(X))$ consists of the classes of (-2)-curves on X (see, for example, Rudakov-Shafarevich [26]). For simplicity, we put

$$Aut(\mathrm{Nef}(X)) := \{\ g \in \mathrm{O}^+(S_X) \ \mid\ \mathrm{Nef}(X)^g = \mathrm{Nef}(X)\ \} = Aut_{\mathrm{O}^+(S_X)}(\mathrm{Nef}(X)).$$

6.1. **Complex $K3$ surfaces.** Suppose that X is defined over $\mathbb{C}$. With the cup-product, the second cohomology group $H := H^2(X,\mathbb{Z})$ is an even unimodular lattice of signature $(3,19)$. Let T_X denote the orthogonal complement of S_X in H. We regard a non-zero holomorphic 2-form ω_X on X as a vector of $T_X \otimes \mathbb{C}$, and put

$$C_X := \{\ g \in \mathrm{O}(T_X) \ \mid\ \omega_X^g = \lambda\, \omega_X \ \text{ for some } \ \lambda \in \mathbb{C}^\times\ \}. \tag{6.1}$$

Since H is unimodular, the subgroup $H/(S_X \oplus T_X)$ of the discriminant group $A_{S_X} \oplus A_{T_X}$ of $S_X \oplus T_X$ is the graph of an isomorphism

$$\delta_X : (A_{S_X}, q_{S_X}) \ \xrightarrow{\sim}\ (A_{T_X}, -q_{T_X})$$

by Nikulin [19, Proposition 1.6.1]. Let $\delta_{X*} : \mathrm{O}(q_{S_X}) \xrightarrow{\sim} \mathrm{O}(q_{T_X})$ denote the isomorphism induced by δ_X. Recall that, for an even lattice L, we have a natural homomorphism $\eta_L : \mathrm{O}(L) \to \mathrm{O}(q_L)$.

Theorem 6.1 (Piatetski-Shapiro and Shafarevich [24]). *Via the natural actions of* $\mathrm{Aut}(X)$ *on the lattices* S_X *and* T_X*, the automorphism group* $\mathrm{Aut}(X)$ *of* X *is identified with*

$$\{\ (g,h) \in Aut(\mathrm{Nef}(X)) \times C_X \ \mid\ \delta_{X*}(\eta_{S_X}(g)) = \eta_{T_X}(h)\ \}.$$

Since $\mathrm{O}(q_{T_X})$ is finite, the subgroup

$$G_X := \{\, g \in \mathrm{O}^+(S_X) \mid \delta_{X*}(\eta_{S_X}(g)) \in \eta_{T_X}(C_X)\,\} \tag{6.2}$$

of $\mathrm{O}^+(S_X)$ has finite index.

Corollary 6.2. *The kernel of the natural homomorphism* $\mathrm{Aut}(X) \to \mathrm{O}(S_X)$ *is isomorphic to* $\mathrm{Ker}(\eta_{T_X}) \cap C_X$*, and its image is equal to* $Aut_{G_X}(\mathrm{Nef}(X))$*.*

Remark 6.3. Let ρ denote the Picard number of X. Then T_X is an even lattice of signature $(2, 20-\rho)$ with the discriminant form isomorphic to $-q_{S_X}$. By Nikulin [19, Theorem 1.10.1], there exists an even lattice of signature $(0, 26-\rho)$ with the discriminant form isomorphic to $-q_{S_X}$. Hence S_X has the property [SG1] with $n = 26$ by Nikulin [19, Proposition 1.6.1].

6.2. **Supersingular $K3$ surfaces in odd characteristics.** Suppose that X is a supersingular $K3$ surface defined over an algebraically closed field k of *odd* characteristic $p > 2$. By Artin [1], we know that A_{S_X} is a p-elementary abelian group of rank 2σ, where σ is a positive integer ≤ 10, which is called the *Artin invariant of* X. The $\mathbb{F}_p$-vector space $S_0 := pS_X^{\vee}/pS_X$ of dimension 2σ has a natural quadratic form

$$Q_0 \quad : \quad px \bmod pS_X \mapsto px^2 \bmod p \quad (x \in S_X^{\vee})$$

that takes values in $\mathbb{F}_p$. We denote by $\mathrm{O}(Q_0)$ the finite group of automorphisms of (S_0, Q_0). We have a natural homomorphism $\mathrm{O}(S_X) \to \mathrm{O}(Q_0)$. We denote by $\varphi : S_0 \otimes k \to S_0 \otimes k$ the map $\mathrm{id}_{S_0} \otimes F_k$, where F_k is the Frobenius map of k. Let $c_{\mathrm{DR}} : S_X \to H^2_{\mathrm{DR}}(X/k)$ denote the Chern class map. Then the kernel $\mathrm{Ker}(\bar{c}_{\mathrm{DR}})$ of the induced homomorphism $\bar{c}_{\mathrm{DR}} : S_X \otimes k \to H^2_{\mathrm{DR}}(X/k)$ is contained in $S_0 \otimes k$. The subspace

$$K := \varphi^{-1}(\mathrm{Ker}(\bar{c}_{\mathrm{DR}}))$$

of $S_0 \otimes k$ is called the *period of* X.

Theorem 6.4 (Ogus [22], [23])**.** *Via the natural action of* $\mathrm{Aut}(X)$ *on the lattice* S_X*, the automorphism group* $\mathrm{Aut}(X)$ *of* X *is identified with*

$$\{\, g \in Aut(\mathrm{Nef}(X)) \mid K^g = K\,\}.$$

Since $\mathrm{O}(Q_0)$ is finite, the subgroup

$$G_X := \{\, g \in \mathrm{O}^+(S_X) \mid K^g = K\,\} \tag{6.3}$$

of $\mathrm{O}^+(S_X)$ has finite index.

Corollary 6.5. *The natural homomorphism* $\mathrm{Aut}(X) \to \mathrm{O}(S_X)$ *is injective, and its image is equal to* $Aut_{G_X}(\mathrm{Nef}(X))$*.*

6.3. **Geometric application of Algorithm 5.14.** Let X be a complex algebraic $K3$ surface or a supersingular $K3$ surface in odd characteristic. Let $S := S_X$ be the Néron-Severi lattice of X, and let $G := G_X$ be the subgroup of $\mathrm{O}^+(S)$ defined by (6.2) or (6.3). Suppose that S and G satisfy [G] in Section 3 and [SG1], [SG2], [SG3] in Section 5. Then we can calculate a finite set of generators of

$$\mathrm{Im}(\mathrm{Aut}(X) \to \mathrm{O}(S_X)) = Aut_G(\mathrm{Nef}(X))$$

by Algorithm 5.14, provided that we find a Weyl vector of an $\mathcal{R}^*_{II|S}$-chamber D_0 contained in the $\mathcal{R}^*_S$-chamber $E = \mathrm{Nef}(X)$.

7. COMPLEX ELLIPTIC $K3$ SURFACES WITH PICARD NUMBER 3

Let X be a complex $K3$ surface with Picard number 3 and with a Jacobian fibration

$$\phi : X \to \mathbb{P}^1$$

whose Mordell-Weil group MW_ϕ is of rank 1. We demonstrate Algorithm 5.14 on X and calculate $\mathrm{Aut}(X)$ under the assumption that the period of X is *generic*, which means that the group C_X defined by (6.1) is equal to $\{\pm 1\}$.

We denote by $f_\phi \in S_X$ the class of a fiber of ϕ and by $z_\phi \in S_X$ the class of the zero section of ϕ. Then there exists $v_3 \in S_X$ such that f_ϕ, z_ϕ, v_3 form a basis of S_X, and that the Gram matrix M of S_X with respect to f_ϕ, z_ϕ, v_3 is

$$M = \begin{bmatrix} 0 & 1 & 0 \\ 1 & -2 & 0 \\ 0 & 0 & -2k \end{bmatrix},$$

where $k := -v_3^2/2$. Since MW_ϕ is of rank 1, there are no reducible fibers of ϕ and hence we have $k > 1$. A vector $\xi f_\phi + \eta z_\phi + \zeta v_3$ of $S_X \otimes \mathbb{R}$ is written as $[\xi, \eta, \zeta]$. The discriminant group A_{S_X} of S_X is a cyclic group of order $2k$ generated by $u_3 \bmod S_X$, where $u_3 := [0, 0, -1/2k]$. Since $k > 1$, we have $-1 \notin \mathrm{Ker}(\eta_{T_X})$. Therefore, by Corollary 6.2 and the assumption $C_X = \{\pm 1\}$, the natural homomorphism $\mathrm{Aut}(X) \to \mathrm{O}^+(S_X)$ is injective, and $\mathrm{Aut}(X) \cong Aut_{G_X}(\mathrm{Nef}(X))$ holds, where G_X is defined by (6.2). We describe the group G_X in terms of matrices. Note that the class $a := [2, 1, 0] \in S_X$ is ample, and hence $g \in \mathrm{O}(S_X)$ belongs to $\mathrm{O}^+(S_X)$ if and only if $\langle a^g, a\rangle_{S_X} > 0$. Therefore G_X is canonically identified with

$$\{\ g \in \mathrm{GL}_3(\mathbb{Z}) \ \mid\ g\, M\, {}^t g = M,\ \ a\, g\, M\, {}^t a > 0,\ \ u_3 g \equiv \pm u_3 \bmod \mathbb{Z}^3\ \}.$$

In particular, the subgroup G_X of $\mathrm{O}^+(S_X)$ satisfies [G] in Section 3.

We have a canonical bijection

$$\mathrm{MW}_\phi \ \cong\ \{\ [ks^2, 1, s] \ \mid\ s \in \mathbb{Z}\ \}$$

by sending sections of ϕ to the classes of their images. The Mordell-Weil group $\mathrm{MW}_\phi \cong \mathbb{Z}$ acts on X as translations. We also have the inversion automorphism $\iota_X : X \to X$ induced by the multiplication by -1 on the generic fiber of ϕ. Therefore $\mathrm{Aut}(X)$ contains $\mathrm{MW}_\phi \rtimes \langle \iota_X \rangle \cong \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z}$, which is generated by the two involutions

$$h_1 := \iota_X = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \quad h_2 := \begin{bmatrix} 1 & 0 & 0 \\ k & 1 & -1 \\ 2k & 0 & -1 \end{bmatrix}.$$

Applying Algorithm 5.14 to $S := S_X$ and $II := \mathrm{II}_{1,9}$, we will obtain $h_3, \dots, h_N \in \mathrm{O}^+(S_X)$ such that $\mathrm{Aut}(X) \cong Aut_{G_X}(\mathrm{Nef}(X))$ is generated by $\{h_1, h_2, h_3, \dots, h_N\}$.

The norm of $[1, x, y] \in S_X \otimes \mathbb{R}$ is $2x - 2x^2 - 2ky^2$. Hence, by the map $[1, x, y] \mapsto (x, y)$, the hyperbolic plane $\mathbb{H}_{S_X}$ associated with S_X is identified with

$$H_X := \{\ (x, y) \in \mathbb{R}^2 \ \mid\ (x - 1/2)^2 + (\sqrt{k}y)^2 < 1/4\ \}.$$

The vector f_ϕ corresponds to the point $(0, 0)$ of $\overline{H}_X$, and the hyperplane $(z_\phi)^\perp$ is given by $x = 1/2$.

Let $\mathcal{D}_0$ be the $\mathcal{R}^*_{II}$-chamber with the Weyl vector $w_0 \in II = U \oplus E_8$ given in Section 4.1. The primitive embedding $i : S_X \hookrightarrow II$ such that $D_0 := i^{-1}(\mathcal{D}_0)$ is



FIGURE 7.1. H_X

an $\mathcal{R}^*_{II|S_X}$-chamber contained in $\mathrm{Nef}(X)$ is obtained by the following method. We choose a primitive vector $u_0 \in E_8$ such that $u_0^2 = -2k$, and let $i_0 : S_X \hookrightarrow II$ be the primitive embedding given by

$$i_0(f_\phi) = f_U, \quad i_0(z_\phi) = z_U, \quad i_0(v_3) = u_0.$$

Let $a' \in S_X$ be a sufficiently general vector in the ample cone of X. (We have used $a' = [2A, A, -1]$ with A large enough.) We calculate the finite set

$$\{\ r \in \mathcal{R}_{II} \ \mid\ \langle w_0, r\rangle_{II} > 0,\ \ \langle i_0(a'), r\rangle_{II} < 0\ \} = \{r_1, \dots, r_N\}$$

by Algorithm 2.3, and sort the elements of this set in such a way that

$$t_1 < t_2 < \cdots < t_N, \quad \text{where} \quad t_i = -\frac{\langle i_0(a'), r_i\rangle_{II}}{\langle w_0, r_i\rangle_{II}}.$$

Then the oriented line segment in $II \otimes \mathbb{R}$ from $i_0(a')$ to w_0 intersects the hyperplanes $(r_1)^\perp, \dots, (r_N)^\perp$ in this order. Let $i : S_X \hookrightarrow II$ be the composite of i_0 with the product $s_{r_1} \cdots s_{r_N}$ of reflections with respect to r_i. Then $i(a') \in \mathcal{D}_0$ holds, and hence $D_0 = i^{-1}(\mathcal{D}_0)$ is an $\mathcal{R}^*_{II|S_X}$-chamber containing a' in its interior. In particular, we have $D_0 \subset \mathrm{Nef}(X)$. Starting from this $\mathcal{R}^*_{II|S_X}$-chamber D_0, we execute Algorithm 5.14.

Remark 7.1. The embedding i depends on the choice of the ample class a'.

Example 7.2. We present the result for the case $-2k = -16$. Elements of II is written in terms of the basis $f_U, z_U, e_1, \dots, e_8$. The embedding i is given by

$$i(f_\phi) = f_U, \quad i(z_\phi) = z_U, \quad i(v_3) = [0, 0, 11, 7, 14, 21, 17, 13, 9, 5].$$

As the elements of $\mathcal{C}$, we obtained nineteen $\mathcal{R}^*_{II|S_X}$-chambers $D_0, \dots, D_{18}$. It turns out that we have $Aut_{G_X}(D_i) = \{1\}$ for $i = 0, \dots, 18$. Their walls and adjacency relation are given in Table 7.1, where (a, b, c) is the wall defined by $a + bx + cy = 0$ in H_X. A (-2)-*wall* means a wall that belongs to $\mathcal{R}^*_{S_X}$. For example, the chamber D_{14} has three walls $-1 - x - 9y = 0, 2x + 5y = 0$ and $1 + 6y = 0$. The chamber adjacent to D_{14} along $-1 - x - 9y = 0$ is D_{12}, the chamber adjacent to D_{14} along

D_0	$(1,-2,0)$	(-2)-wall	D_1	$(1,-1,5)$	D_2	D_2	$(1,-2,0)$	(-2)-wall
	$(0,1,5)$	D_1		$(0,1,4)$	D_3		$(-1,1,-5)$	D_1
	$(0,0,-1)$	$\cong D_0$ by h_1		$(0,-1,-5)$	D_0		$(0,1,4)$	D_4
D_3	$(1,-1,5)$	D_4	D_4	$(1,-1,4)$	D_6	D_5	$(1,-1,5)$	D_7
	$(0,1,3)$	D_5		$(0,1,3)$	D_7		$(0,2,5)$	D_8
	$(0,-1,-4)$	D_1		$(-1,1,-5)$	D_3		$(0,-1,-3)$	D_3
				$(0,-1,-4)$	D_2			
D_6	$(1,-2,0)$	(-2)-wall	D_7	$(0,-1,-3)$	D_4	D_8	$(1,-1,5)$	D_{10}
	$(1,0,7)$	D_9		$(1,0,7)$	D_{11}		$(0,1,2)$	$\cong D_8$ by h_2
	$(-1,1,-4)$	D_4		$(0,2,5)$	D_{10}		$(0,-2,-5)$	D_5
				$(-1,1,-5)$	D_5			
D_9	$(1,-2,0)$	(-2)-wall	D_{10}	$(0,-2,-5)$	D_7	D_{11}	$(1,-1,4)$	D_{12}
	$(-1,0,-7)$	D_6		$(1,0,7)$	D_{13}		$(-1,0,-7)$	D_7
	$(0,1,3)$	D_{12}		$(-1,1,-5)$	D_8		$(0,2,5)$	D_{13}
D_{12}	$(-1,1,-4)$	D_{11}	D_{13}	$(0,-2,-5)$	D_{11}	D_{14}	$(1,0,6)$	$\cong D_{14}$ by h_3
	$(1,1,9)$	D_{14}		$(-1,0,-7)$	D_{10}		$(0,2,5)$	D_{16}
	$(0,-1,-3)$	D_9		$(1,1,9)$	D_{15}		$(-1,-1,-9)$	D_{12}
D_{15}	$(1,-1,4)$	D_{16}	D_{16}	$(0,-2,-5)$	D_{14}	D_{17}	$(1,-1,4)$	D_{18}
	$(-1,-1,-9)$	D_{13}		$(1,2,11)$	D_{18}		$(-1,-2,-11)$	D_{15}
	$(1,2,11)$	D_{17}		$(-1,1,-4)$	D_{15}		$(0,1,2)$	$\cong D_{17}$ by h_2
D_{18}	$(1,0,6)$	$\cong D_{18}$ by h_3						
	$(-1,-2,-11)$	D_{16}						
	$(-1,1,-4)$	D_{17}						

TABLE 7.1. Chambers for the case $-2k=-16$



FIGURE 7.2. F_E for the case $-2k=-16$

$2x+5y=0$ is D_{16}, and the chamber D' adjacent to D_{14} along $1+6y=0$ satisfies $D'=D_{14}^{h_3}$, where

$$h_3 := \begin{bmatrix} 17 & 8 & -3 \\ 0 & 1 & 0 \\ 96 & 48 & -17 \end{bmatrix}.$$

It follows that $\mathrm{Aut}(X)$ is generated by h_1, h_2 and h_3. Note that $h_3 \notin \mathrm{MW}_\phi$, because $f_\phi^{h_3}$ is not equal to f_ϕ. The union F_E of $D_0,\dots,D_{18}$ is given in Figure 7.2, in which the (-2)-walls are depicted by thick lines. Since only one (-2)-wall appears in Figure 7.2, we see that $\mathrm{Aut}(X)$ acts on the set of (-2)-curves in X transitively.

Example 7.3. We present the result for the case $-2k=-22$ in Table 7.2 and Figure 7.3. The embedding i is given by

$$i(f_\phi)=f_U,\quad i(z_\phi)=z_U,\quad i(v_3)=[0,0,12,8,16,24,20,16,11,6].$$

D_0	$(1,-2,0)$	(-2)-wall	D_1	$(0,1,5)$	D_3	D_2	$(1,-2,0)$	(-2)-wall
	$(0,1,6)$	D_1		$(0,-1,-6)$	D_0		$(-1,1,-6)$	D_1
	$(0,0,-1)$	$\cong D_0$ by h_1		$(1,-1,6)$	D_2		$(0,1,5)$	D_4
D_3	$(0,-1,-5)$	D_1	D_4	$(1,-1,5)$	D_6	D_5	$(0,-1,-4)$	D_3
	$(1,-1,6)$	D_4		$(0,1,4)$	D_7		$(1,-1,6)$	D_7
	$(0,1,4)$	D_5		$(-1,1,-6)$	D_3		$(0,1,3)$	D_8
				$(0,-1,-5)$	D_2			
D_6	$(1,-2,0)$	(-2)-wall	D_7	$(1,-1,5)$	D_9	D_8	$(0,-1,-3)$	D_5
	$(-1,1,-5)$	D_4		$(0,1,3)$	D_{10}		$(1,-1,6)$	D_{10}
	$(0,1,4)$	D_9		$(-1,1,-6)$	D_5		$(0,2,5)$	D_{11}
				$(0,-1,-4)$	D_4			
D_9	$(0,-1,-4)$	D_6	D_{10}	$(0,-1,-3)$	D_7	D_{11}	$(0,-2,-5)$	D_8
	$(-1,1,-5)$	D_7		$(1,1,11)$	D_{13}		$(1,1,11)$	D_{14}
	$(1,0,8)$	D_{12}		$(-1,1,-6)$	D_8		$(0,1,2)$	$\cong D_{11}$ by h_2
D_{12}	$(1,-1,4)$	D_{16}	D_{13}	$(1,0,8)$	D_{18}	D_{14}	$(0,1,2)$	$\cong D_{14}$ by h_2
	$(1,1,11)$	D_{15}		$(-1,-1,-11)$	D_{10}		$(-1,-1,-11)$	D_{11}
	$(-1,0,-8)$	D_9		$(0,2,5)$	D_{17}		$(1,-1,6)$	D_{17}
D_{15}	$(1,-1,4)$	D_{19}	D_{16}	$(1,-2,0)$	(-2)-wall	D_{17}	$(0,-2,-5)$	D_{13}
	$(-1,-1,-11)$	D_{12}		$(1,1,11)$	D_{19}		$(-1,1,-6)$	D_{14}
	$(0,1,3)$	D_{20}		$(-1,1,-4)$	D_{12}		$(1,0,8)$	D_{21}
D_{18}	$(1,-1,5)$	D_{20}	D_{19}	$(-1,1,-4)$	D_{15}	D_{20}	$(0,-1,-3)$	D_{15}
	$(0,2,5)$	D_{21}		$(-1,-1,-11)$	D_{16}		$(-1,1,-5)$	D_{18}
	$(-1,0,-8)$	D_{13}		$(2,-1,11)$	D_{22}		$(1,1,10)$	D_{23}
D_{21}	$(0,-2,-5)$	D_{18}	D_{22}	$(1,-2,0)$	(-2)-wall	D_{23}	$(-1,-1,-10)$	D_{20}
	$(-1,0,-8)$	D_{17}		$(-2,1,-11)$	D_{19}		$(2,1,17)$	$\cong D_{25}$ by h_3
	$(1,1,10)$	D_{24}		$(1,0,7)$	D_{25}		$(0,2,5)$	D_{26}
D_{24}	$(1,-1,5)$	D_{26}	D_{25}	$(1,-2,0)$	(-2)-wall	D_{26}	$(0,-2,-5)$	D_{23}
	$(2,3,22)$	(-2)-wall		$(2,1,17)$	$\cong D_{23}$ by h_3		$(2,3,22)$	(-2)-wall
	$(-1,-1,-10)$	D_{21}		$(-1,0,-7)$	D_{22}		$(-1,1,-5)$	D_{24}

TABLE 7.2. Chambers for the case $-2k = -22$



FIGURE 7.3. F_E for the case $-2k = -22$

We have $Aut_{G_X}(D_i) = \{1\}$ for $i = 0, \ldots, 26$. The extra automorphism is

$$h_3 := \begin{bmatrix} 20 & 9 & -3 \\ 7 & 2 & -1 \\ 154 & 66 & -23 \end{bmatrix}.$$

The set of (-2)-curves on X is decomposed into at most two orbits under the action of $\mathrm{Aut}(X)$.

Remark 7.4. Example 1.1 in Introduction is the case $-2k = -24$. The region F_E consists of 46 chambers, and is given in Figure 7.4.

8. OTHER EXAMPLES

We applied Algorithm 5.14 to the following $K3$ surfaces.



Figure 7.4. F_E for the case $-2k = -24$

(1) The complex Fermat quartic surface $X \subset \mathbb{P}^3$. The Picard number of X is 20. Note that X contains 48 lines. We can calculate a Gram matrix of S_X, because S_X is generated by the classes of 20 lines in X. We embed S_X into $II = \mathrm{II}_{1,25}$ primitively in such a way that w_{0,S_X} is equal to the class of a hyperplane section.

(2) The double plane $\pi : X \to \mathbb{P}^2$ branched along the Fermat curve $B \subset \mathbb{P}^2$ of degree 6 in characteristic 5. This $K3$ surface X is supersingular with Artin invariant 1, and contains 252 rational curves that are mapped to lines on $\mathbb{P}^2$ isomorphically by π. The lattice S_X is generated by the classes of 22 curves among them (see [29]). Thus we can calculate a Gram matrix of S_X. We embed S_X into $II = \mathrm{II}_{1,25}$ primitively in such a way that w_{0,S_X} is equal to the class of the pull-back of a line on $\mathbb{P}^2$.

The computation for these two cases did not terminate in a reasonable time, because there are too many G_X-congruence classes of $\mathcal{R}^*_{II|S_X}$-chambers. However, we obtained many interesting automorphisms of these $K3$ surfaces. Detailed study of these automorphisms will appear in other papers.

References

[1] M. Artin. Supersingular $K3$ surfaces. *Ann. Sci. École Norm. Sup. (4)*, 7:543–567 (1975), 1974.

[2] A. Ash, D. Mumford, M. Rapoport, and Yung-Sheng Tai. *Smooth compactifications of locally symmetric varieties.* Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 2010.

[3] R. E. Borcherds. The Leech lattice. *Proc. Roy. Soc. London Ser. A*, 398(1815):365–376, 1985.

[4] R. E. Borcherds. Automorphism groups of Lorentzian lattices. *J. Algebra*, 111(1):133–153, 1987.

[5] R. E. Borcherds. Coxeter groups, Lorentzian lattices, and $K3$ surfaces. *Internat. Math. Res. Notices*, (19):1011–1031, 1998.

[6] V. Chvátal. *Linear programming.* A Series of Books in the Mathematical Sciences. W. H. Freeman and Company, New York, 1983.

[7] J. H. Conway. The automorphism group of the 26-dimensional even unimodular Lorentzian lattice. *J. Algebra*, 80(1):159–163, 1983.

[8] J. H. Conway, R. A. Parker, and N. J. A. Sloane. The covering radius of the Leech lattice. *Proc. Roy. Soc. London Ser. A*, 380(1779):261–290, 1982.

[9] J. H. Conway and N. J. A. Sloane. Lorentzian forms for the Leech lattice. *Bull. Amer. Math. Soc. (N.S.)*, 6(2):215–217, 1982.

[10] J. H. Conway and N. J. A. Sloane. Twenty-three constructions for the Leech lattice. *Proc. Roy. Soc. London Ser. A*, 381(1781):275–283, 1982.

[11] J. H. Conway and N. J. A. Sloane. *Sphere packings, lattices and groups*, volume 290 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, New York, third edition, 1999.

[12] I. Dolgachev and S. Kondō. A supersingular $K3$ surface in characteristic 2 and the Leech lattice. *Int. Math. Res. Not.*, (1):1–23, 2003.

[13] I. Dolgachev and Jonghae Keum. Birational automorphisms of quartic Hessian surfaces. *Trans. Amer. Math. Soc.*, 354(8):3031–3057 (electronic), 2002.

[14] T. Granlund et al. Gmp, the gnu multiple precision arithmetic library. Available from http://gmplib.org/index.html.

[15] J. Keum and S. Kondō. The automorphism groups of Kummer surfaces associated with the product of two elliptic curves. *Trans. Amer. Math. Soc.*, 353(4):1469–1487 (electronic), 2001.

[16] S. Kondō. The automorphism group of a generic Jacobian Kummer surface. *J. Algebraic Geom.*, 7(3):589–609, 1998.

[17] S. Kondō and I. Shimada. The automorphism group of a supersingular $K3$ surface with Artin invariant 1 in characteristic 3, 2012. arXiv:1205.6520, to appear in Int. Math. Res. Not.

[18] M. Lieblich and D. Maulik. A note on the cone conjecture for $K3$ surfaces in positive characteristic. preprint, 2011, arXiv:1102.3377.

[19] V. V. Nikulin. Integer symmetric bilinear forms and some of their geometric applications. *Izv. Akad. Nauk SSSR Ser. Mat.*, 43(1):111–177, 238, 1979. English translation: Math USSR-Izv. 14 (1979), no. 1, 103–167 (1980).

[20] V. V. Nikulin. Quotient-groups of groups of automorphisms of hyperbolic forms by subgroups generated by 2-reflections. Algebro-geometric applications. In *Current problems in mathematics, Vol. 18*, pages 3–114. Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow, 1981.

[21] V. V. Nikulin. On the classification of hyperbolic root systems of rank three. *Tr. Mat. Inst. Steklova*, 230:256, 2000.

[22] A. Ogus. Supersingular $K3$ crystals. In *Journées de Géométrie Algébrique de Rennes (Rennes, 1978), Vol. II*, volume 64 of *Astérisque*, pages 3–86. Soc. Math. France, Paris, 1979.

[23] A. Ogus. A crystalline Torelli theorem for supersingular $K3$ surfaces. In *Arithmetic and geometry, Vol. II*, volume 36 of *Progr. Math.*, pages 361–394. Birkhäuser Boston, Boston, MA, 1983.

[24] I. I. Pjateckiĭ-Šapiro and I. R. Šafarevič. Torelli's theorem for algebraic surfaces of type K3. *Izv. Akad. Nauk SSSR Ser. Mat.*, 35:530–572, 1971. Reprinted in I. R. Shafarevich, Collected Mathematical Papers, Springer-Verlag, Berlin, 1989, pp. 516–557.

[25] J. G. Ratcliffe. *Foundations of hyperbolic manifolds*, volume 149 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994.

[26] A. N. Rudakov and I. R. Shafarevich. Surfaces of type $K3$ over fields of finite characteristic. In *Current problems in mathematics, Vol. 18*, pages 115–207. Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow, 1981. Reprinted in I. R. Shafarevich, Collected Mathematical Papers, Springer-Verlag, Berlin, 1989, pp. 657–714.

[27] J.-P. Serre. *A course in arithmetic*. Springer-Verlag, New York, 1973. Translated from the French, Graduate Texts in Mathematics, No. 7.

[28] I. Shimada. Supersingular $K3$ surfaces in odd characteristic and sextic double planes. *Math. Ann.*, 328(3):451–468, 2004.

[29] I. Shimada. Projective models of the supersingular $K3$ surface with Artin invariant 1 in characteristic 5, 2012. preprint, arXiv:1201.4533.

[30] H. Sterk. Finiteness results for algebraic $K3$ surfaces. *Math. Z.*, 189(4):507–513, 1985.

[31] È. B. Vinberg. The two most algebraic $K3$ surfaces. *Math. Ann.*, 265(1):1–21, 1983.

[32] È. B. Vinberg. Classification of 2-reflective hyperbolic lattices of rank 4. *Tr. Mosk. Mat. Obs.*, 68:44–76, 2007.

Department of Mathematics, Graduate School of Science, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima, 739-8526 JAPAN

E-mail address: `shimada@math.sci.hiroshima-u.ac.jp`